

Key Indicators for Haryana from NFHS-3	NFHS-3 (2005-06)	Residence		Education				NFHS-2 (1998-99)	NFHS-1 (1992-93)
		Urban	Rural	No education ⁴	<5 years complete ⁴	5-9 years complete ⁴	10 years complete and above ⁴		
Marriage and Fertility									
1. Women age 20-24 married by age 18 (%)	41.2	33.3	44.8	65.0	*	52.7	19.7	41.5	57.3
2. Men age 25-29 married by age 21 (%)	33.7	30.0	35.8	*	*	(41.4)	20.3	na	na
3. Total fertility rate (children per woman)	2.7	2.2	2.9	3.4	2.7	2.5	2.3	2.9	4.0
4. Women age 15-19 who were already mothers or pregnant at the time of the survey (%)	12.1	3.7	15.2	25.9	*	13.9	6.5	na	na
5. Median age at first birth for women age 25-49	20.3	20.9	20.0	19.5	20.8	19.9	22.3	20.1	19.8
6. Married women with 2 living children wanting no more children ¹ (%)	88.7	87.9	89.1	84.7	*	88.6	92.0	81.1	63.4
6a. Two sons	97.4	96.7	97.7	95.1	*	100.0	98.4	95.2	85.7
6b. One son, one daughter	91.8	92.7	91.4	82.9	*	94.7	95.4	84.8	67.6
6c. Two daughters	(25.8)	*	(17.9)	*	*	*	*	17.6	8.7
Family Planning (currently married women, age 15-49)									
Current use									
7. Any method (%)	63.4	66.5	62.0	66.4	54.7	61.5	61.1	62.4	49.7
8. Any modern method (%)	58.3	59.7	57.7	62.9	53.0	54.3	55.2	53.2	44.4
8a. Female sterilization (%)	38.2	28.2	42.3	53.7	28.6	33.4	16.8	38.7	29.8
8b. Male sterilization (%)	0.7	1.3	0.5	0.7	3.8	0.7	0.4	2.1	5.0
8c. IUD (%)	4.7	5.6	4.4	1.8	6.4	5.1	9.3	3.6	3.2
8d. Pill (%)	2.8	4.2	2.2	1.4	1.6	3.5	4.5	2.1	1.2
8e. Condom (%)	11.8	20.3	8.2	5.2	12.5	11.5	23.6	6.8	5.2
Unmet need for family planning									
9. Total unmet need (%)	8.3	6.9	8.9	6.4	4.8	10.9	9.2	7.6	16.4
9a. For spacing (%)	3.1	2.0	3.6	1.7	3.2	4.1	4.5	2.9	8.8
9b. For limiting (%)	5.2	4.9	5.3	4.7	1.6	6.8	4.7	4.7	7.6
Maternal and Child Health									
Maternity care (for births in the last 3 years)									
10. Mothers who had at least 3 antenatal care visits for their last birth (%)	58.8	75.5	52.8	39.9	*	60.6	80.0	38.2	45.8
11. Mothers who consumed IFA for 90 days or more when they were pregnant with their last child (%)	28.3	42.9	23.1	11.4	*	31.1	47.7	na	na
12. Births assisted by a doctor/nurse/LHV/ANM/other health personnel (%) ²	54.2	79.0	45.9	29.6	*	57.9	81.8	42.1	31.5
13. Institutional births (%) ²	39.4	66.7	30.3	14.6	*	41.4	69.6	22.4	17.4
14. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/other health personnel within 2 days of delivery for their last birth (%) ²	40.6	63.3	32.5	19.1	*	38.8	69.5	na	na
Child immunization and vitamin A supplementation²									
15a. Children 12-23 months fully immunized (BCG, measles, and 3 doses each of polio/DPT) (%)	65.3	(82.2)	60.3	45.0	*	73.4	85.9	62.7	53.5
15b. Children 12-23 months who have received BCG (%)	84.9	(88.9)	83.7	66.5	*	94.9	100.0	86.8	77.4
15c. Children 12-23 months who have received 3 doses of polio vaccine (%)	82.8	(88.9)	81.0	72.8	*	84.6	94.2	74.3	67.7
15d. Children 12-23 months who have received 3 doses of DPT vaccine (%)	74.2	(84.4)	71.2	54.8	*	85.5	90.1	71.1	66.8
15e. Children 12-23 months who have received measles vaccine (%)	75.5	(84.4)	72.8	52.4	*	87.3	95.6	72.2	60.9
16. Children age 12-35 months who received a vitamin A dose in last 6 months (%)	16.0	20.0	14.7	8.7	*	16.3	25.0	na	na
Treatment of childhood diseases (children under 3 years)²									
17. Children with diarrhoea in the last 2 weeks who received ORS (%)	24.8	*	16.5	*	*	(23.9)	(43.2)	25.7	7.6
18. Children with diarrhoea in the last 2 weeks taken to a health facility (%)	82.7	*	81.0	*	*	(90.3)	(85.9)	92.6	67.1
19. Children with acute respiratory infection or fever in the last 2 weeks taken to a health facility (%)	88.1	*	87.3	*	*	*	(92.4)	na	na
Child Feeding Practices and Nutritional Status of Children^{2,3}									
20. Children under 3 years breastfed within one hour of birth (%)	22.3	25.2	21.3	19.8	*	22.0	24.4	11.7	2.5
21. Children age 0-5 months exclusively breastfed (%)	16.9	*	16.7	(17.0)	*	(23.5)	(12.5)	na	na
22. Children age 6-9 months receiving solid or semi-solid food and breastmilk (%)	44.8	(42.9)	45.7	(38.6)	*	(35.3)	(59.8)	na	na
23. Children under 3 years who are stunted (%)	43.3	36.1	45.7	52.8	*	45.5	26.9	55.6	50.5
24. Children under 3 years who are wasted (%)	22.4	23.8	22.0	25.3	*	21.8	18.1	7.8	7.8
25. Children under 3 years who are underweight (%)	38.2	36.7	38.7	46.6	*	41.8	23.3	29.9	31.0
Nutritional Status of Ever-Married Adults (age 15-49)									
26. Women whose Body Mass Index is below normal (%)	27.8	16.9	32.5	34.5	35.0	25.3	17.1	25.9	na
27. Men whose Body Mass Index is below normal (%)	26.8	19.3	30.0	39.2	39.8	28.1	16.5	na	na
28. Women who are overweight or obese (%)	21.0	34.8	15.0	13.9	16.8	22.6	33.3	16.6	na
29. Men who are overweight or obese (%)	14.4	26.5	9.3	4.8	9.4	13.2	21.5	na	na
Anaemia among Children and Adults									
30. Children age 6-35 months who are anaemic (%)	82.3	79.8	83.3	88.3	*	80.1	77.0	83.9	na
31. Ever-married women age 15-49 who are anaemic (%)	56.6	55.6	56.9	57.6	60.7	58.7	51.7	47.0	na
32. Pregnant women age 15-49 who are anaemic (%)	69.7	61.5	71.7	80.7	*	66.7	(65.5)	55.5	na
33. Ever-married men age 15-49 who are anaemic (%)	17.6	17.0	17.8	20.8	29.6	17.0	13.9	na	na
Knowledge of HIV/AIDS among Ever-Married Adults (age 15-49)									
34. Women who have heard of AIDS (%)	60.0	76.5	53.1	32.5	50.6	73.7	96.3	44.3	na
35. Men who have heard of AIDS (%)	87.1	91.0	85.5	64.1	79.9	89.9	97.6	na	na
36. Women who know that consistent condom use can reduce the chances of getting HIV/AIDS (%)	45.3	59.5	39.3	19.5	30.5	54.5	84.3	na	na
37. Men who know that consistent condom use can reduce the chances of getting HIV/AIDS (%)	79.9	86.2	77.2	50.8	72.3	81.4	94.7	na	na
Women's Empowerment									
38. Currently married women who usually participate in household decisions (%)	41.7	49.5	38.4	44.3	37.6	39.6	39.8	na	na
39. Ever-married women who have ever experienced spousal violence (%)	27.3	23.9	28.8	36.3	(37.7)	24.1	13.9	na	na

na : Not available

nc : Not calculated because there are no cases

() Based on 25-49 unweighted cases

* Based on fewer than 25 unweighted cases

¹ Excludes pregnant women.

² Based on the last 2 births in the 3 years before the survey to ever-married women.

³ Based on WHO standard.

⁴ For children education refers to mother's education. Children with missing information on the mother's education are not included in the education columns.