



सत्यमेव जयते

Government of India

Ministry of Health and Family Welfare

NATIONAL FAMILY HEALTH SURVEY - 4

2015-16

DISTRICT FACT SHEET LUDHIANA PUNJAB



(स्थापना / Established in 1956)
बेहतर भविष्य के लिए क्षमता निर्माण
Capacity Building for a Better Future

International Institute for Population Sciences
(Deemed University)
Mumbai

Introduction

The National Family Health Survey 2015-16 (NFHS-4), the fourth in the NFHS series, provides information on population, health and nutrition for India and each State / Union territory. NFHS-4, for the first time, provides district-level estimates for many important indicators.

The contents of previous rounds of NFHS are generally retained and additional components are added from one round to another. In this round, information on malaria prevention, migration in the context of HIV, abortion, violence during pregnancy etc. have been added. The scope of clinical, anthropometric, and biochemical testing (CAB) or Biomarker component has been expanded to include measurement of blood pressure and blood glucose levels. NFHS-4 sample has been designed to provide district and higher level estimates of various indicators covered in the survey. However, estimates of indicators of sexual behaviour, husband's background and woman's work, HIV/AIDS knowledge, attitudes and behaviour, and, domestic violence will be available at State and national level only.

As in the earlier rounds, the Ministry of Health and Family Welfare, Government of India designated International Institute for Population Sciences, Mumbai as the nodal agency to conduct NFHS-4. The main objective of each successive round of the NFHS has been to provide essential data on health and family welfare and emerging issues in this area. NFHS-4 data will be useful in setting benchmarks and examining the progress in health sector the country has made over time. Besides providing evidence for the effectiveness of the ongoing programmes, the data from NFHS-4 help in identifying need for new programmes with area specific focus.

Four Survey Schedules - Household, Woman's, Man's and Biomarker - were canvassed in local language using Computer Assisted Personal Interviewing (CAPI). In the Household Schedule, information was collected on all usual members of the household and visitors who stayed in the household the previous night as well as socio-economic characteristics of the household, water and sanitation, health insurance, number of deaths in the household in the three years preceding the survey etc. Information on the woman's characteristics, marriage, fertility, children's immunizations and childcare, nutrition, contraception, reproductive health, sexual behaviour, HIV/AIDS, domestic violence, etc. was canvassed in the Woman's Schedule. The Man's Schedule covered the man's characteristics, marriage, his number of children, contraception, fertility preferences, nutrition, sexual behaviour, attitudes towards gender roles, HIV/AIDS, etc. The Biomarker Schedule covered measurements of height, weight and haemoglobin levels for children; measurements of height, weight, haemoglobin levels, blood pressure, and random blood glucose level for women aged 15-49 years and men aged 15-54 years. In addition, women and men were requested to provide a few drops of blood from a finger prick for laboratory testing for HIV.

This fact sheet provides information on key indicators and trends for Ludhiana. NFHS-4 fieldwork for Punjab was conducted from 28 January 2016 to 20 June 2016 by Society for Promotion of Youth & Masses (SPYM). In Ludhiana, information was gathered from 814 households, 918 women, and 129 men. The fact sheet shows information for urban and rural areas and the district as a whole because the percent urban population in Ludhiana is between 30-70%, which provides a sufficiently large sample to produce reliable estimates of most indicators for both urban and rural areas.

Ludhiana, Punjab - Key Indicators

Indicators	NFHS-4 (2015-16)		
	Urban	Rural	Total
Population and Household Profile			
1. Population (female) age 6 years and above who ever attended school (%)	83.9	79.1	82.0
2. Population below age 15 years (%)	24.5	20.7	23.0
3. Sex ratio of the total population (females per 1,000 males)	823	929	862
4. Sex ratio at birth for children born in the last five years (females per 1,000 males)	751	979	827
5. Children under age 5 years whose birth was registered (%)	97.3	100.0	98.2
6. Households with electricity (%)	100.0	99.7	99.9
7. Households with an improved drinking-water source ¹ (%)	100.0	100.0	100.0
8. Households using improved sanitation facility ² (%)	75.3	90.1	80.8
9. Households using clean fuel for cooking ³ (%)	92.4	50.1	76.8
10. Households using iodized salt (%)	98.4	99.0	98.6
11. Households with any usual member covered by a health scheme or health insurance (%)	18.4	33.5	24.0
Characteristics of Adults (age 15-49)			
12. Women who are literate (%)	83.9	86.6	85.0
13. Men who are literate (%)	89.3	83.3	87.2
14. Women with 10 or more years of schooling (%)	67.9	54.0	62.5
Marriage and Fertility			
15. Women age 20-24 years married before age 18 years (%)	3.3	10.2	6.4
16. Men age 25-29 years married before age 21 years (%)	*	*	*
17. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	(2.4)	1.5	2.0
Current Use of Family Planning Methods (currently married women age 15-49 years)			
18. Any method ⁴ (%)	72.3	73.5	72.8
19. Any modern method ⁴ (%)	60.1	64.3	61.7
20. Female sterilization (%)	30.4	40.7	34.2
21. Male sterilization (%)	0.2	1.3	0.6
22. IUD/PPIUD (%)	9.2	3.8	7.2
23. Pill (%)	2.1	1.2	1.8
24. Condom (%)	18.3	17.4	18.0
Unmet Need for Family Planning (currently married women age 15-49 years)⁵			
25. Total unmet need (%)	4.5	5.9	5.0
26. Unmet need for spacing (%)	1.5	3.1	2.1
Quality of Family Planning Services			
27. Health worker ever talked to female non-users about family planning (%)	18.5	19.0	18.7
28. Current users ever told about side effects of current method ⁶ (%)	(63.5)	(56.5)	60.9

¹ Piped water into dwelling/yard/plot, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rainwater, community RO plant.

² Flush to piped sewer system, flush to septic tank, flush to pit latrine, ventilated improved pit (VIP)/biogas latrine, pit latrine with slab, twin pit/composting toilet, which is not shared with any other household. ³ Electricity, LPG/natural gas, biogas. ⁴ Includes other methods that are not shown separately

⁵ Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.
- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

Women who are classified as infertile have no unmet need because they are not at risk of becoming pregnant. Unmet need for family planning is the sum of unmet need for spacing plus unmet need for limiting.

⁶ Based on current users of female sterilization, IUD/PPIUD, injectables and pill who started using that method in the past 5 years.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

Ludhiana, Punjab - Key Indicators

Indicators	NFHS-4 (2015-16)		
	Urban	Rural	Total
Maternal and Child Health			
Maternity Care (for last birth in the 5 years before the survey)			
29. Mothers who had antenatal check-up in the first trimester (%)	75.7	86.8	79.9
30. Mothers who had at least 4 antenatal care visits (%)	67.6	73.8	69.9
31. Mothers whose last birth was protected against neonatal tetanus ⁷ (%)	98.9	97.1	98.2
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	41.5	42.6	42.0
33. Mothers who had full antenatal care ⁸ (%)	29.3	32.9	30.6
34. Registered pregnancies for which the mother received Mother and Child Protection (MCP) card (%)	93.0	96.5	94.4
35. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery (%)	80.5	89.3	83.8
36. Mothers who received financial assistance under Janani Suraksha Yojana (JSY) for births delivered in an institution (%)	6.1	18.7	11.2
37. Average out of pocket expenditure per delivery in public health facility (Rs.)	(1,004)	1,481	1,269
38. Children born at home who were taken to a health facility for check-up within 24 hours of birth (%)	*	*	(2.7)
39. Children who received a health check after birth from a doctor/nurse/LHV/ANM/ midwife/other health personnel within 2 days of birth (%)	43.1	56.3	48.1
Delivery Care (for births in the 5 years before the survey)			
40. Institutional births (%)	78.5	89.5	82.5
41. Institutional births in public facility (%)	26.2	51.3	35.4
42. Home delivery conducted by skilled health personnel (out of total deliveries) (%)	9.2	4.5	7.5
43. Births assisted by a doctor/nurse/LHV/ANM/other health personnel (%)	85.5	94.0	88.6
44. Births delivered by caesarean section (%)	23.4	20.3	22.3
45. Births in a private health facility delivered by caesarean section (%)	38.2	32.0	36.3
46. Births in a public health facility delivered by caesarean section (%)	(13.2)	15.8	14.6
Child Immunizations and Vitamin A Supplementation			
47. Children age 12-23 months fully immunized (BCG, measles, and 3 doses each of polio and DPT) (%)	*	*	(72.3)
48. Children age 12-23 months who have received BCG (%)	*	*	(94.6)
49. Children age 12-23 months who have received 3 doses of polio vaccine (%)	*	*	(81.6)
50. Children age 12-23 months who have received 3 doses of DPT vaccine (%)	*	*	(88.1)
51. Children age 12-23 months who have received measles vaccine (%)	*	*	(82.5)
52. Children age 12-23 months who have received 3 doses of Hepatitis B vaccine (%)	*	*	(81.9)
53. Children age 9-59 months who received a vitamin A dose in last 6 months (%)	47.8	43.3	46.2
54. Children age 12-23 months who received most of the vaccinations in public health facility (%)	*	*	(80.2)
55. Children age 12-23 months who received most of the vaccinations in private health facility (%)	*	*	(19.9)
Treatment of Childhood Diseases (children under age 5 years)			
56. Prevalence of diarrhoea (reported) in the last 2 weeks preceding the survey (%)	4.3	3.9	4.2
57. Children with diarrhoea in the last 2 weeks who received oral rehydration salts (ORS) (%)	*	*	*
58. Children with diarrhoea in the last 2 weeks who received zinc (%)	*	*	*
59. Children with diarrhoea in the last 2 weeks taken to a health facility (%)	*	*	*
60. Prevalence of symptoms of acute respiratory infection (ARI) in the last 2 weeks preceding the survey (%)	2.7	0.0	1.7
61. Children with fever or symptoms of ARI in the last 2 weeks preceding the survey taken to a health facility (%)	*	*	*
Child Feeding Practices and Nutritional Status of Children			
62. Children under age 3 years breastfed within one hour of birth ⁹ (%)	30.8	36.4	32.9
63. Children under age 6 months exclusively breastfed ¹⁰ (%)	*	*	(61.3)
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	*	*	*
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10,11} (%)	*	(6.6)	(9.0)
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10,11} (%)	*	*	*
67. Total children age 6-23 months receiving an adequate diet ^{10,11} (%)	*	(5.6)	6.6
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	35.1	21.3	30.1
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	16.9	19.6	17.8
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	1.8	5.5	3.1
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	30.6	24.7	28.5

⁷ Includes mothers with two injections during the pregnancy of her last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth. Not exactly comparable with NFHS-3 due to differences in definition. ⁸ Full antenatal care is at least four antenatal visits, at least one tetanus toxoid (TT) injection and iron folic acid tablets or syrup taken for 100 or more days. ⁹ Based on the last child born in the 5 years before the survey. ¹⁰ Based on the youngest child living with the mother. ¹¹ Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group). ¹² Below -2 standard deviations, based on the WHO standard. ¹³ Below -3 standard deviations, based on the WHO standard.

Ludhiana, Punjab - Key Indicators

Indicators	NFHS-4 (2015-16)		
	Urban	Rural	Total
Nutritional Status of Adults (age 15-49 years)			
72. Women whose Body Mass Index (BMI) is below normal (BMI < 18.5 kg/m ²) ¹⁴ (%)	8.6	10.2	9.2
73. Men whose Body Mass Index (BMI) is below normal (BMI < 18.5 kg/m ²) (%)	9.4	12.1	10.3
74. Women who are overweight or obese (BMI ≥ 25.0 kg/m ²) ¹⁴ (%)	33.8	38.1	35.5
75. Men who are overweight or obese (BMI ≥ 25.0 kg/m ²) (%)	32.7	23.3	29.4
Anaemia among Children and Adults¹⁵			
76. Children age 6-59 months who are anaemic (<11.0 g/dl) (%)	62.9	57.1	60.8
77. Non-pregnant women age 15-49 years who are anaemic (<12.0 g/dl) (%)	66.0	66.0	66.0
78. Pregnant women age 15-49 years who are anaemic (<11.0 g/dl) (%)	*	*	(45.6)
79. All women age 15-49 years who are anaemic (%)	64.7	66.0	65.2
80. Men age 15-49 years who are anaemic (<13.0 g/dl) (%)	29.5	37.0	32.0
Blood Sugar Level among Adults (age 15-49 years)¹⁶			
Women			
81. Blood sugar level - high (>140 mg/dl) (%)	4.5	6.6	5.3
82. Blood sugar level - very high (>160 mg/dl) (%)	1.9	3.3	2.4
Men			
83. Blood sugar level - high (>140 mg/dl) (%)	6.6	7.3	6.8
84. Blood sugar level - very high (>160 mg/dl) (%)	4.1	0.0	2.7
Hypertension among Adults (age 15-49 years)			
Women			
85. Slightly above normal (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	10.3	8.5	9.6
86. Moderately high (Systolic 160-179 mm of Hg and/or Diastolic 100-109 mm of Hg) (%)	1.2	2.6	1.7
87. Very high (Systolic ≥180 mm of Hg and/or Diastolic ≥110 mm of Hg) (%)	0.0	1.1	0.4
Men			
88. Slightly above normal (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	13.6	21.9	16.5
89. Moderately high (Systolic 160-179 mm of Hg and/or Diastolic 100-109 mm of Hg) (%)	3.2	1.4	2.6
90. Very high (Systolic ≥180 mm of Hg and/or Diastolic ≥110 mm of Hg) (%)	0.0	6.3	2.2
Women Age 15-49 Years Who Have Ever Undergone Examinations of:			
91. Cervix (%)	29.8	36.3	32.4
92. Breast (%)	21.8	24.9	23.0
93. Oral cavity (%)	50.4	46.2	48.8

¹⁴ Excludes pregnant women and women with a birth in the preceding 2 months. ¹⁵ Haemoglobin in grams per decilitre (g/dl). Among children, prevalence is adjusted for altitude. Among adults, prevalence is adjusted for altitude and for smoking status. ¹⁶ Random blood sugar measurement (including those under medication).

INTERNATIONAL INSTITUTE FOR POPULATION SCIENCES

Vision: “To position IIPS as a premier teaching and research institution in population sciences responsive to emerging national and global needs based on values of inclusion, sensitivity and rights protection.”

Mission: “The Institute will strive to be a centre of excellence on population, health and development issues through high quality education, teaching and research. This will be achieved by (a) creating competent professionals, (b) generating and disseminating scientific knowledge and evidence, (c) collaboration and exchange of knowledge, and (d) advocacy and awareness.”

For additional information, please contact:

Director/Project Coordinator (NFHS-4)
International Institute for Population Sciences
Govandi Station Road, Deonar
Mumbai - 400 088 (India)
Telephone: 022-4237 2442
Fax: 022-25563257
Email: nfhs42013@gmail.com, director@iips.net
Website: <http://www.rchiips.org/nfhs>
<http://www.iipsindia.org>

Chief Director (Stat.)
Ministry of Health and Family Welfare
Government of India
Nirman Bhavan
New Delhi 110 108
Telephone: 011 – 23062288
Fax: 011 - 23062288
Email: cdstat@nic.in
Website: <http://www.mohfw.nic.in>
<https://www.nrhm-mis.nic.in>

Director (Stat.)
Ministry of Health and Family Welfare
Government of India
Nirman Bhavan
New Delhi 110 108
Telephone: 011 - 23062647
Fax: 011 - 23062647
Email: jdstat2-mohfw@nic.in
Website: <http://www.mohfw.nic.in>
<https://www.nrhm-mis.nic.in>

Technical assistance for NFHS-4 was provided by the USAID-supported DHS project at ICF, and assistance for the HIV components was provided by NACO and NARI. Funding assistance was provided by Ministry of Health and Family Welfare, Government of India and:



USAID
FROM THE AMERICAN PEOPLE



MacArthur
Foundation

The opinions in this publication do not necessarily reflect the views of the funding agencies.

For additional information on NFHS-4, visit <http://www.rchiips.org/nfhs>

For related information, visit <http://www.iipsindia.org> or <http://www.mohfw.nic.in>