



सत्यमेव जयते
Government of India

Ministry of Health and Family Welfare

Compendium of Fact Sheets

KEY INDICATORS STATE AND DISTRICTS OF MEGHALAYA

National Family
Health Survey (NFHS-5)

2019-20



(स्थापना / Established in 1956)

बेहतर भविष्य के लिए क्षमता निर्माण
Capacity Building for a Better Future

International Institute for Population Sciences
(Deemed University)

Suggested citation: International Institute for Population Sciences (IIPS) and ICF. 2020.
National Family Health Survey (NFHS)-5, *State and District Factsheets*, Meghalaya. Mumbai: IIPS.

CONTRIBUTORS

Sarang Pedgaonkar
H. Lhungdim
K. S. James
Nirbhay Kumar Singh

© *International Institute for Population Sciences, Mumbai*

For additional information about the 2019-20 National Family Health Survey (NFHS-5), please contact:

International Institute for Population Sciences, Govandi Station Road, Deonar, Mumbai-400 088

Telephone: 022-4237 2467

E-mail: nfhs52017@gmail.com; director@iips.net

For related information, visit <http://www.rchiips.org/nfhs> or <http://www.iipsindia.ac.in>

Key Indicators Content

Content	Page No.
State	
Meghalaya	1
District	
1. East Garo Hills	7
2. East Jantia Hills	13
3. East Khasi Hills	19
4. North Garo Hills	25
5. Ribhoi	31
6. South Garo Hills	37
7. South West Garo Hills	43
8. South West Khasi Hills	49
9. West Garo Hills	55
10. West Jaintia Hills	61
11. West Khasi Hills	67



सत्यमेव जयते
Government of India

Ministry of Health and Family Welfare

NATIONAL FAMILY HEALTH SURVEY - 5

2019-20

STATE FACT SHEET

MEGHALAYA



(स्थापना / Established in 1956)

बेहतर भविष्य के लिए क्षमता निर्माण
Capacity Building for a Better Future

International Institute for Population Sciences
(Deemed University)

Introduction

The National Family Health Survey 2019-20 (NFHS-5), the fifth in the NFHS series, provides information on population, health, and nutrition for India and each state/union territory (UT). Like NFHS-4, NFHS-5 also provides district-level estimates for many important indicators.

The contents of NFHS-5 are similar to NFHS-4 to allow comparisons over time. However, NFHS-5 includes some new topics, such as preschool education, disability, access to a toilet facility, death registration, bathing practices during menstruation, and methods and reasons for abortion. The scope of clinical, anthropometric, and biochemical testing (CAB) has also been expanded to include measurement of waist and hip circumferences, and the age range for the measurement of blood pressure and blood glucose has been expanded. However, HIV testing has been dropped. The NFHS-5 sample has been designed to provide national, state/union territory (UT), and district level estimates of various indicators covered in the survey. However, estimates of indicators of sexual behaviour; husband's background and woman's work; HIV/AIDS knowledge, attitudes and behaviour; and domestic violence are available only at the state/union territory (UT) and national level.

As in the earlier rounds, the Ministry of Health and Family Welfare, Government of India, designated the International Institute for Population Sciences, Mumbai, as the nodal agency to conduct NFHS-5. The main objective of each successive round of the NFHS has been to provide high-quality data on health and family welfare and emerging issues in this area. NFHS-5 data will be useful in setting benchmarks and examining the progress the health sector has made over time. Besides providing evidence for the effectiveness of ongoing programmes, the data from NFHS-5 help in identifying the need for new programmes with an area specific focus and identifying groups that are most in need of essential services.

Four Survey Schedules - Household, Woman's, Man's, and Biomarker - were canvassed in local languages using Computer Assisted Personal Interviewing (CAPI). In the Household Schedule, information was collected on all usual members of the household and visitors who stayed in the household the previous night, as well as socio-economic characteristics of the household; water, sanitation, and hygiene; health insurance coverage; disabilities; land ownership; number of deaths in the household in the three years preceding the survey; and the ownership and use of mosquito nets. The Woman's Schedule covered a wide variety of topics, including the woman's characteristics, marriage, fertility, contraception, children's immunizations and healthcare, nutrition, reproductive health, sexual behaviour, HIV/AIDS, women's empowerment, and domestic violence. The Man's Schedule covered the man's characteristics, marriage, his number of children, contraception, fertility preferences, nutrition, sexual behaviour, health issues, attitudes towards gender roles, and HIV/AIDS. The Biomarker Schedule covered measurements of height, weight, and haemoglobin levels for children; measurements of height, weight, waist and hip circumference, and haemoglobin levels for women age 15-49 years and men age 15-54 years; and blood pressure and random blood glucose levels for women and men age 15 years and over. In addition, women and men were requested to provide a few additional drops of blood from a finger prick for laboratory testing for HbA1c, malaria parasites, and Vitamin D3.

Readers should be cautious while interpreting and comparing the trends as some States/UTs may have smaller sample size. Moreover, at the time of survey, Ayushman Bharat AB-PMJAY and Pradhan Mantri Surakshit Matritva Abhiyan (PMSMA) were not fully rolled out and hence, their coverage may not have been factored in the results of indicator 12 (percentage of households with any usual member covered under a health insurance/financing scheme) and indicator 41 (percentage of mothers who received 4 or more antenatal care check-ups).

This fact sheet provides information on key indicators and trends for Meghalaya. NFHS-5 fieldwork for Meghalaya was conducted from 8 July, 2019 to 15 November, 2019 by IQVIA Consulting and Information Services India Pvt. Ltd. Information was gathered from 10,148 households, 13,089 women, and 1,824 men. Fact sheets for each district in Meghalaya are also available separately.

Meghalaya - Key Indicators

Indicators	NFHS-5 (2019-20)			NFHS-4 (2015-16)
	Urban	Rural	Total	Total
Population and Household Profile				
1. Female population age 6 years and above who ever attended school (%)	94.7	83.4	85.9	83.0
2. Population below age 15 years (%)	29.7	39.2	37.3	36.5
3. Sex ratio of the total population (females per 1,000 males)	1,118	1,020	1,039	1,005
4. Sex ratio at birth for children born in the last five years (females per 1,000 males)	915	1,001	989	1,009
5. Children under age 5 years whose birth was registered with the civil authority (%)	87.5	81.2	82.1	79.8
6. Deaths in the last 3 years registered with the civil authority (%)	60.9	52.8	54.4	na
7. Population living in households with electricity (%)	97.3	90.6	91.9	91.8
8. Population living in households with an improved drinking-water source ¹ (%)	93.6	75.6	79.2	70.3
9. Population living in households that use an improved sanitation facility ² (%)	81.3	83.3	82.9	61.4
10. Households using clean fuel for cooking ³ (%)	76.7	21.7	33.7	21.8
11. Households using iodized salt (%)	97.5	88.7	90.6	99.1
12. Households with any usual member covered under a health insurance/financing scheme (%)	52.8	66.5	63.5	34.6
13. Children age 5 years who attended pre-primary school during the school year 2019-20 (%)	36.3	30.3	31.2	na
Characteristics of Adults (age 15-49 years)				
14. Women who are literate ⁴ (%)	97.1	85.5	88.2	na
15. Men who are literate ⁴ (%)	92.9	81.5	83.7	na
16. Women with 10 or more years of schooling (%)	61.4	27.3	35.1	33.6
17. Men with 10 or more years of schooling (%)	63.9	27.7	34.7	34.5
18. Women who have ever used the internet (%)	57.8	28.0	34.7	na
19. Men who have ever used the internet (%)	59.2	38.5	42.1	na
Marriage and Fertility				
20. Women age 20-24 years married before age 18 years (%)	9.1	19.1	16.9	16.9
21. Men age 25-29 years married before age 21 years (%)	(4.8)	20.8	17.9	13.6
22. Total fertility rate (children per woman)	1.6	3.3	2.9	3.0
23. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	3.2	8.4	7.2	8.6
24. Adolescent fertility rate for women age 15-19 years ⁵	18	58	49	48
Infant and Child Mortality Rates (per 1,000 live births)				
25. Neonatal mortality rate (NNMR)	(14.2)	20.6	19.8	18.3
26. Infant mortality rate (IMR)	(23.4)	33.6	32.3	29.9
27. Under-five mortality rate (U5MR)	23.4	42.6	40.0	39.6
Current Use of Family Planning Methods (currently married women age 15-49 years)				
28. Any method ⁶ (%)	25.9	27.8	27.4	24.3
29. Any modern method ⁶ (%)	21.0	22.9	22.5	21.9
30. Female sterilization (%)	7.1	5.2	5.6	6.2
31. Male sterilization (%)	0.0	0.0	0.0	0.0
32. IUD/PPIUD (%)	2.9	4.7	4.4	2.1
33. Pill (%)	6.0	8.9	8.3	11.7
34. Condom (%)	3.9	2.4	2.7	1.3
35. Injectables (%)	0.7	1.2	1.1	0.6
Unmet Need for Family Planning (currently married women age 15-49 years)				
36. Total unmet need ⁷ (%)	21.9	28.2	26.9	21.2
37. Unmet need for spacing ⁷ (%)	13.3	19.5	18.3	15.3
Quality of Family Planning Services				
38. Health worker ever talked to female non-users about family planning (%)	18.7	30.4	27.4	24.2
39. Current users ever told about side effects of current method ⁸ (%)	61.4	68.4	67.2	61.3

Note: Major indicators are highlighted in grey.

LHV = Lady health visitor; ANM = Auxiliary nurse midwife; na = Not available

* Percentage not shown; based on fewer than 25 unweighted cases

For all indicators other than 25, 26, 27: () Based on 25-49 unweighted cases

For indicators 25, 26 and 27: () Based on 250-499 unweighted person-years of exposure to the risk of death

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rainwater, tanker truck, cart with small tank, bottled water, community RO plant.

²Flush to piped sewer system, flush to septic tank, flush to pit latrine, flush to don't know where, ventilated improved pit (VIP)/biogas latrine, pit latrine with slab, twin pit/composting toilet, which is not shared with any other household. This indicator does not denote access to toilet facility.

³Electricity, LPG/natural gas, biogas.

⁴Refers to women/men who completed standard 9 or higher and women/men who can read a whole sentence or part of a sentence.

⁵Equivalent to the age-specific fertility rate for the 3-year period preceding the survey, expressed in terms of births per 1,000 women age 15-19.

⁶Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

⁷Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.

- Pregnant with an unwanted pregnancy.

- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

Women who are classified as infertile have no unmet need because they are not at risk of becoming pregnant. Unmet need for family planning is the sum of unmet need for spacing plus unmet need for limiting.

⁸Based on current users of female sterilization, IUD/PPIUD, injectables, and pills who started using that method in the past 5 years.

Meghalaya - Key Indicators

Indicators	NFHS-5 (2019-20)			NFHS-4 (2015-16)
	Urban	Rural	Total	Total
Maternal and Child Health				
Maternity Care (for last birth in the 5 years before the survey)				
40. Mothers who had an antenatal check-up in the first trimester (%)	63.2	52.3	53.9	53.3
41. Mothers who had at least 4 antenatal care visits (%)	67.5	49.6	52.2	50.0
42. Mothers whose last birth was protected against neonatal tetanus ⁹ (%)	84.8	81.6	82.1	79.2
43. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	47.5	42.3	43.1	36.2
44. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	28.7	19.2	20.6	13.0
45. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	87.5	94.3	93.3	93.6
46. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery (%)	54.8	42.0	43.9	47.5
47. Average out-of-pocket expenditure per delivery in a public health facility (Rs.)	3,353	3,190	3,219	3,319
48. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	0.0	1.6	1.6	1.4
49. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery (%)	59.1	42.5	44.9	na
Delivery Care (for births in the 5 years before the survey)				
50. Institutional births (%)	82.7	54.3	58.1	51.4
51. Institutional births in public facility (%)	57.4	47.9	49.1	39.5
52. Home births that were conducted by skilled health personnel ¹⁰ (%)	2.2	7.3	6.6	2.6
53. Births attended by skilled health personnel ¹⁰ (%)	82.4	61.2	64.0	53.8
54. Births delivered by caesarean section (%)	21.6	6.1	8.2	7.6
55. Births in a private health facility that were delivered by caesarean section (%)	51.0	34.6	40.8	31.4
56. Births in a public health facility that were delivered by caesarean section (%)	15.2	8.1	9.2	9.8
Child Vaccinations and Vitamin A Supplementation				
57. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ¹¹ (%)	58.3	64.7	63.8	61.4
58. Children age 12-23 months fully vaccinated based on information from vaccination card only ¹² (%)	76.0	80.5	80.0	81.3
59. Children age 12-23 months who have received BCG (%)	93.1	88.7	89.3	85.9
60. Children age 12-23 months who have received 3 doses of polio vaccine ¹³ (%)	63.5	69.3	68.5	71.0
61. Children age 12-23 months who have received 3 doses of penta or DPT vaccine (%)	69.6	73.7	73.1	73.9
62. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	70.4	72.8	72.5	71.8
63. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	12.2	14.5	14.2	na
64. Children age 12-23 months who have received 3 doses of rotavirus vaccine ¹⁴ (%)	5.5	3.9	4.1	na
65. Children age 12-23 months who have received 3 doses of penta or hepatitis B vaccine (%)	56.0	67.3	65.7	62.8
66. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	69.3	59.7	61.0	57.8
67. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	78.3	96.1	93.6	92.4
68. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	17.0	1.3	3.5	4.9
Treatment of Childhood Diseases (children under age 5 years)				
69. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	11.8	10.2	10.4	10.6
70. Children with diarrhoea in the 2 weeks preceding the survey who received oral rehydration salts (ORS) (%)	60.9	75.5	73.2	77.4
71. Children with diarrhoea in the 2 weeks preceding the survey who received zinc (%)	42.2	40.2	40.5	57.8
72. Children with diarrhoea in the 2 weeks preceding the survey taken to a health facility or health provider (%)	69.1	69.2	69.2	69.9
73. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	5.8	4.7	4.8	5.8
74. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	66.6	74.0	72.9	74.9

⁹Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

¹⁰Doctor/nurse/LHV/ANM/midwife/other health personnel.

¹¹Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

¹²Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

¹³Not including polio vaccination given at birth.

¹⁴Since rotavirus is not being provided across all states and districts, the levels should not be compared.

Meghalaya - Key Indicators

Indicators	NFHS-5 (2019-20)			NFHS-4 (2015-16)
	Urban	Rural	Total	Total
Child Feeding Practices and Nutritional Status of Children				
75. Children under age 3 years breastfed within one hour of birth ¹⁵ (%)	70.8	79.9	78.8	60.6
76. Children under age 6 months exclusively breastfed ¹⁶ (%)	(64.5)	39.2	42.7	35.8
77. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁶ (%)	*	66.2	66.8	67.4
78. Breastfeeding children age 6-23 months receiving an adequate diet ^{16, 17} (%)	37.5	28.0	29.0	24.2
79. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{16, 17} (%)	(43.7)	30.1	32.5	19.5
80. Total children age 6-23 months receiving an adequate diet ^{16, 17} (%)	39.5	28.4	29.8	23.5
81. Children under 5 years who are stunted (height-for-age) ¹⁸ (%)	35.1	48.2	46.5	43.8
82. Children under 5 years who are wasted (weight-for-height) ¹⁸ (%)	13.0	12.0	12.1	15.3
83. Children under 5 years who are severely wasted (weight-for-height) ¹⁹ (%)	4.6	4.7	4.7	6.5
84. Children under 5 years who are underweight (weight-for-age) ¹⁸ (%)	22.2	27.3	26.6	28.9
85. Children under 5 years who are overweight (weight-for-height) ²⁰ (%)	4.2	4.0	4.0	3.9
Nutritional Status of Adults (age 15-49 years)				
86. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ²¹ (%)	10.2	11.0	10.8	12.1
87. Men whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) (%)	8.6	9.1	9.0	11.6
88. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ²¹ (%)	17.9	9.7	11.5	12.2
89. Men who are overweight or obese (BMI ≥25.0 kg/m ²) (%)	30.2	10.6	13.9	10.1
90. Women who have high risk waist-to-hip ratio (≥0.85) (%)	55.7	61.9	60.6	na
91. Men who have high risk waist-to-hip ratio (≥0.90) (%)	28.5	24.0	24.7	na
Anaemia among Children and Adults				
92. Children age 6-59 months who are anaemic (<11.0 g/dl) ²² (%)	38.8	46.0	45.1	48.0
93. Non-pregnant women age 15-49 years who are anaemic (<12.0 g/dl) ²² (%)	52.4	54.9	54.4	56.4
94. Pregnant women age 15-49 years who are anaemic (<11.0 g/dl) ²² (%)	40.2	45.9	45.0	53.3
95. All women age 15-49 years who are anaemic ²² (%)	51.8	54.3	53.8	56.2
96. All women age 15-19 years who are anaemic ²² (%)	44.6	54.6	52.5	52.1
97. Men age 15-49 years who are anaemic (<13.0 g/dl) ²² (%)	16.1	27.4	25.5	32.4
98. Men age 15-19 years who are anaemic (<13.0 g/dl) ²² (%)	(7.4)	35.0	30.1	25.2
Blood Sugar Level among Adults (age 15 years and above)				
Women				
99. Blood sugar level - high (141-160 mg/dl) ²³ (%)	3.0	5.5	5.0	na
100. Blood sugar level - very high (>160 mg/dl) ²³ (%)	4.8	2.5	3.0	na
101. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ²³ (%)	10.3	9.3	9.5	na
Men				
102. Blood sugar level - high (141-160 mg/dl) ²³ (%)	6.6	9.0	8.6	na
103. Blood sugar level - very high (>160 mg/dl) ²³ (%)	7.6	3.5	4.2	na
104. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ²³ (%)	16.0	13.4	13.9	na
Hypertension among Adults (age 15 years and above)				
Women				
105. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	13.6	9.0	10.0	na
106. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	4.9	3.6	3.9	na
107. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	24.6	17.1	18.7	na
Men				
108. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	17.8	13.3	14.2	na
109. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	5.8	3.4	3.8	na
110. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	28.5	19.9	21.4	na

¹⁵Based on the last child born in the 3 years before the survey.

¹⁶Based on the youngest child living with the mother.

¹⁷Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹⁸Below -2 standard deviations, based on the WHO standard.

¹⁹Below -3 standard deviations, based on the WHO standard.

²⁰Above +2 standard deviations, based on the WHO standard.

²¹Excludes pregnant women and women with a birth in the preceding 2 months.

²²Haemoglobin in grams per decilitre (g/dl). Among children, prevalence is adjusted for altitude. Among adults, prevalence is adjusted for altitude and for smoking status, if known. As NFHS uses the capillary blood for estimation of anaemia, the results of NFHS-5 need not be compared with other surveys using venous blood.

²³Random blood sugar measurement.

Meghalaya - Key Indicators

Indicators	NFHS-5 (2019-20)			NFHS-4 (2015-16)
	Urban	Rural	Total	Total
Screening for Cancer among Adults (age 30-49 years)				
Women				
111. Ever undergone a screening test for cervical cancer (%)	0.6	0.6	0.6	na
112. Ever undergone a breast examination for breast cancer (%)	1.3	0.2	0.5	na
113. Ever undergone an oral cavity examination for oral cancer (%)	0.4	0.4	0.4	na
Men				
114. Ever undergone an oral cavity examination for oral cancer (%)	2.3	0.9	1.2	na
Knowledge of HIV/AIDS among Adults (age 15-49 years)				
115. Women who have comprehensive knowledge ²⁴ of HIV/AIDS (%)	23.8	11.8	14.5	13.3
116. Men who have comprehensive knowledge ²⁴ of HIV/AIDS (%)	32.5	11.9	15.9	14.2
117. Women who know that consistent condom use can reduce the chance of getting HIV/AIDS (%)	60.6	52.2	54.1	47.0
118. Men who know that consistent condom use can reduce the chance of getting HIV/AIDS (%)	70.6	60.8	62.7	52.1
Women's Empowerment (women age 15-49 years)				
119. Currently married women who usually participate in three household decisions ²⁵ (%)	93.3	92.0	92.3	91.4
120. Women who worked in the last 12 months and were paid in cash (%)	43.1	39.1	40.0	35.9
121. Women owning a house and/or land (alone or jointly with others) (%)	47.8	70.1	65.0	57.3
122. Women having a bank or savings account that they themselves use (%)	78.0	68.2	70.4	54.4
123. Women having a mobile phone that they themselves use (%)	78.2	64.3	67.5	64.3
124. Women age 15-24 years who use hygienic methods of protection during their menstrual period ²⁶ (%)	85.0	59.1	64.9	63.7
Gender Based Violence (age 18-49 years)				
125. Ever-married women age 18-49 years who have ever experienced spousal violence ²⁷ (%)	23.2	14.2	16.0	28.8
126. Ever-married women age 18-49 years who have experienced physical violence during any pregnancy (%)	2.5	1.4	1.6	0.4
127. Young women age 18-29 years who experienced sexual violence by age 18 (%)	0.0	0.6	0.5	0.0
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)				
128. Women age 15 years and above who use any kind of tobacco (%)	27.2	28.5	28.2	na
129. Men age 15 years and above who use any kind of tobacco (%)	47.0	60.6	57.7	na
130. Women age 15 years and above who consume alcohol (%)	1.0	1.6	1.5	na
131. Men age 15 years and above who consume alcohol (%)	28.5	33.5	32.4	na

²⁴Comprehensive knowledge means knowing that consistent use of condoms every time they have sex and having just one uninfected faithful sex partner can reduce the chance of getting HIV/AIDS, knowing that a healthy-looking person can have HIV/AIDS, and rejecting two common misconceptions about transmission or prevention of HIV/AIDS.

²⁵Decisions about health care for herself, making major household purchases, and visits to her family or relatives.

²⁶Locally prepared napkins, sanitary napkins, tampons, and menstrual cups are considered to be hygienic methods of protection.

²⁷Spousal violence is defined as physical and/or sexual violence.



सत्यमेव जयते
Government of India

Ministry of Health and Family Welfare

NATIONAL FAMILY HEALTH SURVEY - 5

2019-20

DISTRICT FACT SHEET

EAST GARO HILLS MEGHALAYA



(स्थापना / Established in 1956)

बेहतर भविष्य के लिए क्षमता निर्माण
Capacity Building for a Better Future

International Institute for Population Sciences
(Deemed University)

Introduction

The National Family Health Survey 2019-20 (NFHS-5), the fifth in the NFHS series, provides information on population, health, and nutrition for India and each state/union territory (UT). Like NFHS-4, NFHS-5 also provides district-level estimates for many important indicators.

The contents of NFHS-5 are similar to NFHS-4 to allow comparisons over time. However, NFHS-5 includes some new topics, such as preschool education, disability, access to a toilet facility, death registration, bathing practices during menstruation, and methods and reasons for abortion. The scope of clinical, anthropometric, and biochemical testing (CAB) has also been expanded to include measurement of waist and hip circumferences, and the age range for the measurement of blood pressure and blood glucose has been expanded. However, HIV testing has been dropped. The NFHS-5 sample has been designed to provide national, state/union territory (UT), and district level estimates of various indicators covered in the survey. However, estimates of indicators of sexual behaviour; husband's background and woman's work; HIV/AIDS knowledge, attitudes and behaviour; and domestic violence are available only at the state/union territory (UT) and national level.

As in the earlier rounds, the Ministry of Health and Family Welfare, Government of India, designated the International Institute for Population Sciences, Mumbai, as the nodal agency to conduct NFHS-5. The main objective of each successive round of the NFHS has been to provide high-quality data on health and family welfare and emerging issues in this area. NFHS-5 data will be useful in setting benchmarks and examining the progress the health sector has made over time. Besides providing evidence for the effectiveness of ongoing programmes, the data from NFHS-5 help in identifying the need for new programmes with an area specific focus and identifying groups that are most in need of essential services.

Four Survey Schedules - Household, Woman's, Man's, and Biomarker - were canvassed in local languages using Computer Assisted Personal Interviewing (CAPI). In the Household Schedule, information was collected on all usual members of the household and visitors who stayed in the household the previous night, as well as socio-economic characteristics of the household; water, sanitation, and hygiene; health insurance coverage; disabilities; land ownership; number of deaths in the household in the three years preceding the survey; and the ownership and use of mosquito nets. The Woman's Schedule covered a wide variety of topics, including the woman's characteristics, marriage, fertility, contraception, children's immunizations and healthcare, nutrition, reproductive health, sexual behaviour, HIV/AIDS, women's empowerment, and domestic violence. The Man's Schedule covered the man's characteristics, marriage, his number of children, contraception, fertility preferences, nutrition, sexual behaviour, health issues, attitudes towards gender roles, and HIV/AIDS. The Biomarker Schedule covered measurements of height, weight, and haemoglobin levels for children; measurements of height, weight, waist and hip circumference, and haemoglobin levels for women age 15-49 years and men age 15-54 years; and blood pressure and random blood glucose levels for women and men age 15 years and over. In addition, women and men were requested to provide a few additional drops of blood from a finger prick for laboratory testing for HbA1c, malaria parasites, and Vitamin D3.

Readers should be cautious while interpreting and comparing the trends as some States/UTs may have smaller sample size. Moreover, at the time of survey, Ayushman Bharat AB-PMJAY and Pradhan Mantri Surakshit Matritva Abhiyan (PMSMA) were not fully rolled out and hence, their coverage may not have been factored in the results of indicator 12 (percentage of households with any usual member covered under a health insurance/financing scheme) and indicator 33 (percentage of mothers who received 4 or more antenatal care check-ups).

This fact sheet provides information on key indicators for East Garo Hills. NFHS-5 fieldwork for Meghalaya was conducted from 8 July, 2019 to 15 November, 2019 by IQVIA Consulting and Information Services India Pvt. Ltd. In East Garo Hills, information was gathered from 919 households, 1,280 women, and 198 men.

East Garo Hills, Meghalaya - Key Indicators

Indicators	NFHS-5 (2019-20)
Population and Household Profile	
	Total
1. Female population age 6 years and above who ever attended school (%)	90.9
2. Population below age 15 years (%)	31.1
3. Sex ratio of the total population (females per 1,000 males)	1,026
4. Sex ratio at birth for children born in the last five years (females per 1,000 males)	1,427
5. Children under age 5 years whose birth was registered with the civil authority (%)	81.4
6. Deaths in the last 3 years registered with the civil authority (%)	45.9
7. Population living in households with electricity (%)	97.4
8. Population living in households with an improved drinking-water source ¹ (%)	67.0
9. Population living in households that use an improved sanitation facility ² (%)	83.9
10. Households using clean fuel for cooking ³ (%)	29.1
11. Households using iodized salt (%)	69.0
12. Households with any usual member covered under a health insurance/financing scheme (%)	71.9
13. Children age 5 years who attended pre-primary school during the school year 2019-20 (%)	3.2
Characteristics of Women (age 15-49 years)	
14. Women who are literate ⁴ (%)	86.7
15. Women with 10 or more years of schooling (%)	40.5
Marriage and Fertility	
16. Women age 20-24 years married before age 18 years (%)	13.7
17. Births in the 5 years preceding the survey that are third or higher order (%)	1.9
18. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	2.2
19. Women age 15-24 years who use hygienic methods of protection during their menstrual period ⁵ (%)	75.7
Current Use of Family Planning Methods (currently married women age 15-49 years)	
20. Any method ⁶ (%)	37.0
21. Any modern method ⁶ (%)	35.8
22. Female sterilization (%)	4.4
23. Male sterilization (%)	0.0
24. IUD/PPIUD (%)	4.7
25. Pill (%)	18.5
26. Condom (%)	4.9
27. Injectables (%)	2.2
Unmet Need for Family Planning (currently married women age 15-49 years)	
28. Total unmet need ⁷ (%)	21.0
29. Unmet need for spacing ⁷ (%)	12.4
Quality of Family Planning Services	
30. Health worker ever talked to female non-users about family planning (%)	22.1
31. Current users ever told about side effects of current method ⁸ (%)	79.2

Note: Indicator estimates for NFHS-4 are not shown in this table since no comparable estimates are available from NFHS-4 in this district due to district boundary changes or a newly formed district. Major indicators are highlighted in grey.

LHV = Lady health visitor, ANM = Auxiliary nurse midwife

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rainwater, tanker truck, cart with small tank, bottled water, community RO plant.

²Flush to piped sewer system, flush to septic tank, flush to pit latrine, flush to don't know where, ventilated improved pit (VIP)/biogas latrine, pit latrine with slab, twin pit/composting toilet, which is not shared with any other household. This indicator does not denote access to toilet facility completely.

³Electricity, LPG/natural gas, biogas.

⁴Refers to women who completed standard 9 or higher and women who can read a whole sentence or part of a sentence.

⁵Locally prepared napkins, sanitary napkins, tampons, and menstrual cups are considered to be hygienic methods of protection.

⁶Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

⁷Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.
- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

Women who are classified as infecund have no unmet need because they are not at risk of becoming pregnant. Unmet need for family planning is the sum of unmet need for spacing plus unmet need for limiting.

⁸Based on current users of female sterilization, IUD/PPIUD, injectables, and pills who started using that method in the past 5 years.

East Garo Hills, Meghalaya - Key Indicators

Indicators	NFHS-5 (2019-20)
Maternal and Child Health	
Maternity Care (for last birth in the 5 years before the survey)	
32. Mothers who had an antenatal check-up in the first trimester (%)	59.2
33. Mothers who had at least 4 antenatal care visits (%)	25.5
34. Mothers whose last birth was protected against neonatal tetanus ⁹ (%)	74.5
35. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	28.8
36. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	7.0
37. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	92.1
38. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery (%)	59.9
39. Average out-of-pocket expenditure per delivery in a public health facility (Rs.)	2,473
40. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	0.0
41. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery (%)	57.5
Delivery Care (for births in the 5 years before the survey)	
42. Institutional births (%)	73.8
43. Institutional births in public facility (%)	73.5
44. Home births that were conducted by skilled health personnel ¹⁰ (%)	7.5
45. Births attended by skilled health personnel ¹⁰ (%)	80.8
46. Births delivered by caesarean section (%)	4.0
47. Births in a private health facility that were delivered by caesarean section (%)	*
48. Births in a public health facility that were delivered by caesarean section (%)	5.0
Child Vaccinations and Vitamin A Supplementation	
49. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ¹¹ (%)	72.4
50. Children age 12-23 months fully vaccinated based on information from vaccination card only ¹² (%)	(94.5)
51. Children age 12-23 months who have received BCG (%)	81.5
52. Children age 12-23 months who have received 3 doses of polio vaccine ¹³ (%)	76.6
53. Children age 12-23 months who have received 3 doses of penta or DPT vaccine (%)	83.3
54. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	83.3
55. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	19.3
56. Children age 12-23 months who have received 3 doses of rotavirus vaccine ¹⁴ (%)	12.5
57. Children age 12-23 months who have received 3 doses of penta or hepatitis B vaccine (%)	74.8
58. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	56.6
59. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	(93.8)
60. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	(0.0)
Treatment of Childhood Diseases (children under age 5 years)	
61. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	9.2
62. Children with diarrhoea in the 2 weeks preceding the survey who received oral rehydration salts (ORS) (%)	(79.1)
63. Children with diarrhoea in the 2 weeks preceding the survey who received zinc (%)	(63.3)
64. Children with diarrhoea in the 2 weeks preceding the survey taken to a health facility or health provider (%)	(83.3)
65. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	1.6
66. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	64.1

⁹Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

¹⁰Doctor/nurse/LHV/ANM/midwife/other health personnel.

¹¹Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

¹²Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

¹³Not including polio vaccination given at birth.

¹⁴Since rotavirus is not being provided across all states and districts, the levels should not be compared.

East Garo Hills, Meghalaya - Key Indicators

Indicators	NFHS-5 (2019-20)
Child Feeding Practices and Nutritional Status of Children	
Total	
67. Children under age 3 years breastfed within one hour of birth ¹⁵ (%)	48.0
68. Children under age 6 months exclusively breastfed ¹⁶ (%)	*
69. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁶ (%)	*
70. Breastfeeding children age 6-23 months receiving an adequate diet ^{16, 17} (%)	41.3
71. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{16, 17} (%)	*
72. Total children age 6-23 months receiving an adequate diet ^{16, 17} (%)	43.9
73. Children under 5 years who are stunted (height-for-age) ¹⁸ (%)	39.7
74. Children under 5 years who are wasted (weight-for-height) ¹⁸ (%)	20.1
75. Children under 5 years who are severely wasted (weight-for-height) ¹⁹ (%)	6.2
76. Children under 5 years who are underweight (weight-for-age) ¹⁸ (%)	26.5
77. Children under 5 years who are overweight (weight-for-height) ²⁰ (%)	4.1
Nutritional Status of Women (age 15-49 years)	
78. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ²¹ (%)	8.5
79. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ²¹ (%)	7.0
80. Women who have high risk waist-to-hip ratio (≥0.85) (%)	63.7
Anaemia among Children and Women	
81. Children age 6-59 months who are anaemic (<11.0 g/dl) ²² (%)	28.0
82. Non-pregnant women age 15-49 years who are anaemic (<12.0 g/dl) ²² (%)	52.1
83. Pregnant women age 15-49 years who are anaemic (<11.0 g/dl) ²² (%)	29.1
84. All women age 15-49 years who are anaemic ²² (%)	50.9
85. All women age 15-19 years who are anaemic ²² (%)	55.7
Blood Sugar Level among Adults (age 15 years and above)	
Women	
86. Blood sugar level - high (141-160 mg/dl) ²³ (%)	6.3
87. Blood sugar level - very high (>160 mg/dl) ²³ (%)	2.0
88. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ²³ (%)	9.3
Men	
89. Blood sugar level - high (141-160 mg/dl) ²³ (%)	12.0
90. Blood sugar level - very high (>160 mg/dl) ²³ (%)	3.6
91. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ²³ (%)	16.2
Hypertension among Adults (age 15 years and above)	
Women	
92. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	8.5
93. Moderately or severely elevated blood pressure (Systolic ≥160mm of Hg and/or Diastolic ≥100mm of Hg) (%)	2.9
94. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	15.3
Men	
95. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	16.4
96. Moderately or severely elevated blood pressure (Systolic ≥160mm of Hg and/or Diastolic ≥100mm of Hg) (%)	4.4
97. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	23.2
Screening for Cancer among Women (age 30-49 years)	
98. Ever undergone a screening test for cervical cancer (%)	0.2
99. Ever undergone a breast examination for breast cancer (%)	0.2
100. Ever undergone an oral cavity examination for oral cancer (%)	1.1
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)	
101. Women age 15 years and above who use any kind of tobacco (%)	6.3
102. Men age 15 years and above who use any kind of tobacco (%)	54.0
103. Women age 15 years and above who consume alcohol (%)	0.9
104. Men age 15 years and above who consume alcohol (%)	36.8

¹⁵Based on the last child born in the 3 years before the survey.

¹⁶Based on the youngest child living with the mother.

¹⁷Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹⁸Below -2 standard deviations, based on the WHO standard.

¹⁹Below -3 standard deviations, based on the WHO standard.

²⁰Above +2 standard deviations, based on the WHO standard.

²¹Excludes pregnant women and women with a birth in the preceding 2 months.

²²Haemoglobin in grams per decilitre (g/dl). Among children, prevalence is adjusted for altitude. Among women, prevalence is adjusted for altitude and for smoking status, if known. As NFHS uses the capillary blood for estimation of anaemia, the results of NFHS-5 need not be compared with other surveys using venous blood.

²³Random blood sugar measurement.

NOTES



सत्यमेव जयते
Government of India

Ministry of Health and Family Welfare

NATIONAL FAMILY HEALTH SURVEY - 5

2019-20

DISTRICT FACT SHEET

EAST JAIANTIA HILLS

MEGHALAYA



(स्थापना / Established in 1956)

बेहतर भविष्य के लिए क्षमता निर्माण
Capacity Building for a Better Future

International Institute for Population Sciences
(Deemed University)

Introduction

The National Family Health Survey 2019-20 (NFHS-5), the fifth in the NFHS series, provides information on population, health, and nutrition for India and each state/union territory (UT). Like NFHS-4, NFHS-5 also provides district-level estimates for many important indicators.

The contents of NFHS-5 are similar to NFHS-4 to allow comparisons over time. However, NFHS-5 includes some new topics, such as preschool education, disability, access to a toilet facility, death registration, bathing practices during menstruation, and methods and reasons for abortion. The scope of clinical, anthropometric, and biochemical testing (CAB) has also been expanded to include measurement of waist and hip circumferences, and the age range for the measurement of blood pressure and blood glucose has been expanded. However, HIV testing has been dropped. The NFHS-5 sample has been designed to provide national, state/union territory (UT), and district level estimates of various indicators covered in the survey. However, estimates of indicators of sexual behaviour; husband's background and woman's work; HIV/AIDS knowledge, attitudes and behaviour; and domestic violence are available only at the state/union territory (UT) and national level.

As in the earlier rounds, the Ministry of Health and Family Welfare, Government of India, designated the International Institute for Population Sciences, Mumbai, as the nodal agency to conduct NFHS-5. The main objective of each successive round of the NFHS has been to provide high-quality data on health and family welfare and emerging issues in this area. NFHS-5 data will be useful in setting benchmarks and examining the progress the health sector has made over time. Besides providing evidence for the effectiveness of ongoing programmes, the data from NFHS-5 help in identifying the need for new programmes with an area specific focus and identifying groups that are most in need of essential services.

Four Survey Schedules - Household, Woman's, Man's, and Biomarker - were canvassed in local languages using Computer Assisted Personal Interviewing (CAPI). In the Household Schedule, information was collected on all usual members of the household and visitors who stayed in the household the previous night, as well as socio-economic characteristics of the household; water, sanitation, and hygiene; health insurance coverage; disabilities; land ownership; number of deaths in the household in the three years preceding the survey; and the ownership and use of mosquito nets. The Woman's Schedule covered a wide variety of topics, including the woman's characteristics, marriage, fertility, contraception, children's immunizations and healthcare, nutrition, reproductive health, sexual behaviour, HIV/AIDS, women's empowerment, and domestic violence. The Man's Schedule covered the man's characteristics, marriage, his number of children, contraception, fertility preferences, nutrition, sexual behaviour, health issues, attitudes towards gender roles, and HIV/AIDS. The Biomarker Schedule covered measurements of height, weight, and haemoglobin levels for children; measurements of height, weight, waist and hip circumference, and haemoglobin levels for women age 15-49 years and men age 15-54 years; and blood pressure and random blood glucose levels for women and men age 15 years and over. In addition, women and men were requested to provide a few additional drops of blood from a finger prick for laboratory testing for HbA1c, malaria parasites, and Vitamin D3.

Readers should be cautious while interpreting and comparing the trends as some States/UTs may have smaller sample size. Moreover, at the time of survey, Ayushman Bharat AB-PMJAY and Pradhan Mantri Surakshit Matritva Abhiyan (PMSMA) were not fully rolled out and hence, their coverage may not have been factored in the results of indicator 12 (percentage of households with any usual member covered under a health insurance/financing scheme) and indicator 33 (percentage of mothers who received 4 or more antenatal care check-ups).

This fact sheet provides information on key indicators for East Jaintia Hills. NFHS-5 fieldwork for Meghalaya was conducted from 8 July, 2019 to 15 November, 2019 by IQVIA Consulting and Information Services India Pvt. Ltd. In East Jaintia Hills, information was gathered from 924 households, 1,311 women, and 200 men.

East Jaintia Hills, Meghalaya - Key Indicators

Indicators	NFHS-5 (2019-20)
Population and Household Profile	
	Total
1. Female population age 6 years and above who ever attended school (%)	76.0
2. Population below age 15 years (%)	46.1
3. Sex ratio of the total population (females per 1,000 males)	1,066
4. Sex ratio at birth for children born in the last five years (females per 1,000 males)	1,015
5. Children under age 5 years whose birth was registered with the civil authority (%)	83.0
6. Deaths in the last 3 years registered with the civil authority (%)	43.5
7. Population living in households with electricity (%)	85.1
8. Population living in households with an improved drinking-water source ¹ (%)	74.6
9. Population living in households that use an improved sanitation facility ² (%)	83.6
10. Households using clean fuel for cooking ³ (%)	26.5
11. Households using iodized salt (%)	97.2
12. Households with any usual member covered under a health insurance/financing scheme (%)	74.1
13. Children age 5 years who attended pre-primary school during the school year 2019-20 (%)	45.6
Characteristics of Women (age 15-49 years)	
14. Women who are literate ⁴ (%)	76.0
15. Women with 10 or more years of schooling (%)	23.0
Marriage and Fertility	
16. Women age 20-24 years married before age 18 years (%)	25.3
17. Births in the 5 years preceding the survey that are third or higher order (%)	6.3
18. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	14.4
19. Women age 15-24 years who use hygienic methods of protection during their menstrual period ⁵ (%)	47.9
Current Use of Family Planning Methods (currently married women age 15-49 years)	
20. Any method ⁶ (%)	33.6
21. Any modern method ⁶ (%)	24.0
22. Female sterilization (%)	9.2
23. Male sterilization (%)	0.0
24. IUD/PPIUD (%)	1.9
25. Pill (%)	7.6
26. Condom (%)	2.0
27. Injectables (%)	3.4
Unmet Need for Family Planning (currently married women age 15-49 years)	
28. Total unmet need ⁷ (%)	26.0
29. Unmet need for spacing ⁷ (%)	20.1
Quality of Family Planning Services	
30. Health worker ever talked to female non-users about family planning (%)	44.1
31. Current users ever told about side effects of current method ⁸ (%)	81.1

Note: Indicator estimates for NFHS-4 are not shown in this table since no comparable estimates are available from NFHS-4 in this district due to district boundary changes or a newly formed district. Major indicators are highlighted in grey. This indicator does not denote access to toilet facility completely.

LHV = Lady health visitor, ANM = Auxiliary nurse midwife
() Based on 25-49 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rainwater, tanker truck, cart with small tank, bottled water, community RO plant.

²Flush to piped sewer system, flush to septic tank, flush to pit latrine, flush to don't know where, ventilated improved pit (VIP)/biogas latrine, pit latrine with slab, twin pit/composting toilet, which is not shared with any other household.

³Electricity, LPG/natural gas, biogas.

⁴Refers to women who completed standard 9 or higher and women who can read a whole sentence or part of a sentence.

⁵Locally prepared napkins, sanitary napkins, tampons, and menstrual cups are considered to be hygienic methods of protection.

⁶Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

⁷Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
 - Pregnant with a mistimed pregnancy.
 - Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.
- Women are considered to have unmet need for limiting if they are:
- At risk of becoming pregnant, not using contraception, and want no (more) children.
 - Pregnant with an unwanted pregnancy.
 - Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

Women who are classified as infecund have no unmet need because they are not at risk of becoming pregnant. Unmet need for family planning is the sum of unmet need for spacing plus unmet need for limiting.

⁸Based on current users of female sterilization, IUD/PPIUD, injectables, and pills who started using that method in the past 5 years.

East Jaintia Hills, Meghalaya - Key Indicators

Indicators	NFHS-5 (2019-20)
Maternal and Child Health	Total
Maternity Care (for last birth in the 5 years before the survey)	
32. Mothers who had an antenatal check-up in the first trimester (%)	55.5
33. Mothers who had at least 4 antenatal care visits (%)	58.6
34. Mothers whose last birth was protected against neonatal tetanus ⁹ (%)	87.7
35. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	51.9
36. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	25.7
37. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	94.6
38. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery (%)	49.1
39. Average out-of-pocket expenditure per delivery in a public health facility (Rs.)	3,464
40. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	0.4
41. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery (%)	45.0
Delivery Care (for births in the 5 years before the survey)	
42. Institutional births (%)	48.4
43. Institutional births in public facility (%)	36.3
44. Home births that were conducted by skilled health personnel ¹⁰ (%)	6.8
45. Births attended by skilled health personnel ¹⁰ (%)	55.7
46. Births delivered by caesarean section (%)	8.0
47. Births in a private health facility that were delivered by caesarean section (%)	40.3
48. Births in a public health facility that were delivered by caesarean section (%)	8.6
Child Vaccinations and Vitamin A Supplementation	
49. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ¹¹ (%)	74.4
50. Children age 12-23 months fully vaccinated based on information from vaccination card only ¹² (%)	82.4
51. Children age 12-23 months who have received BCG (%)	92.6
52. Children age 12-23 months who have received 3 doses of polio vaccine ¹³ (%)	80.8
53. Children age 12-23 months who have received 3 doses of penta or DPT vaccine (%)	84.0
54. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	83.8
55. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	21.6
56. Children age 12-23 months who have received 3 doses of rotavirus vaccine ¹⁴ (%)	0.6
57. Children age 12-23 months who have received 3 doses of penta or hepatitis B vaccine (%)	77.8
58. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	70.0
59. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	96.9
60. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	1.2
Treatment of Childhood Diseases (children under age 5 years)	
61. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	10.6
62. Children with diarrhoea in the 2 weeks preceding the survey who received oral rehydration salts (ORS) (%)	72.8
63. Children with diarrhoea in the 2 weeks preceding the survey who received zinc (%)	42.2
64. Children with diarrhoea in the 2 weeks preceding the survey taken to a health facility or health provider (%)	57.5
65. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	4.8
66. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	72.0

⁹Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

¹⁰Doctor/nurse/LHV/ANM/midwife/other health personnel.

¹¹Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

¹²Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

¹³Not including polio vaccination given at birth.

¹⁴Since rotavirus is not being provided across all states and districts, the levels should not be compared.

East Jaintia Hills, Meghalaya - Key Indicators

Indicators	NFHS-5 (2019-20)
Child Feeding Practices and Nutritional Status of Children	
Total	
67. Children under age 3 years breastfed within one hour of birth ¹⁵ (%)	85.3
68. Children under age 6 months exclusively breastfed ¹⁶ (%)	50.5
69. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁶ (%)	(63.1)
70. Breastfeeding children age 6-23 months receiving an adequate diet ^{16, 17} (%)	25.4
71. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{16, 17} (%)	25.6
72. Total children age 6-23 months receiving an adequate diet ^{16, 17} (%)	25.5
73. Children under 5 years who are stunted (height-for-age) ¹⁸ (%)	49.8
74. Children under 5 years who are wasted (weight-for-height) ¹⁸ (%)	8.5
75. Children under 5 years who are severely wasted (weight-for-height) ¹⁹ (%)	2.9
76. Children under 5 years who are underweight (weight-for-age) ¹⁸ (%)	23.6
77. Children under 5 years who are overweight (weight-for-height) ²⁰ (%)	3.7
Nutritional Status of Women (age 15-49 years)	
78. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ²¹ (%)	13.5
79. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ²¹ (%)	15.2
80. Women who have high risk waist-to-hip ratio (≥0.85) (%)	62.3
Anaemia among Children and Women	
81. Children age 6-59 months who are anaemic (<11.0 g/dl) ²² (%)	47.2
82. Non-pregnant women age 15-49 years who are anaemic (<12.0 g/dl) ²² (%)	43.7
83. Pregnant women age 15-49 years who are anaemic (<11.0 g/dl) ²² (%)	46.5
84. All women age 15-49 years who are anaemic ²² (%)	43.9
85. All women age 15-19 years who are anaemic ²² (%)	40.6
Blood Sugar Level among Adults (age 15 years and above)	
Women	
86. Blood sugar level - high (141-160 mg/dl) ²³ (%)	3.2
87. Blood sugar level - very high (>160 mg/dl) ²³ (%)	2.1
88. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ²³ (%)	6.3
Men	
89. Blood sugar level - high (141-160 mg/dl) ²³ (%)	4.0
90. Blood sugar level - very high (>160 mg/dl) ²³ (%)	3.3
91. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ²³ (%)	7.7
Hypertension among Adults (age 15 years and above)	
Women	
92. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	9.1
93. Moderately or severely elevated blood pressure (Systolic ≥160mm of Hg and/or Diastolic ≥100mm of Hg) (%)	5.9
94. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	18.7
Men	
95. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	11.1
96. Moderately or severely elevated blood pressure (Systolic ≥160mm of Hg and/or Diastolic ≥100mm of Hg) (%)	4.3
97. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	17.2
Screening for Cancer among Women (age 30-49 years)	
98. Ever undergone a screening test for cervical cancer (%)	0.7
99. Ever undergone a breast examination for breast cancer (%)	0.0
100. Ever undergone an oral cavity examination for oral cancer (%)	0.6
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)	
101. Women age 15 years and above who use any kind of tobacco (%)	49.2
102. Men age 15 years and above who use any kind of tobacco (%)	68.5
103. Women age 15 years and above who consume alcohol (%)	3.6
104. Men age 15 years and above who consume alcohol (%)	42.0

¹⁵Based on the last child born in the 3 years before the survey.

¹⁶Based on the youngest child living with the mother.

¹⁷Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹⁸Below -2 standard deviations, based on the WHO standard.

¹⁹Below -3 standard deviations, based on the WHO standard.

²⁰Above +2 standard deviations, based on the WHO standard.

²¹Excludes pregnant women and women with a birth in the preceding 2 months.

²²Haemoglobin in grams per decilitre (g/dl). Among children, prevalence is adjusted for altitude. Among women, prevalence is adjusted for altitude and for smoking status, if known. As NFHS uses the capillary blood for estimation of anaemia, the results of NFHS-5 need not be compared with other surveys using venous blood.

²³Random blood sugar measurement.

NOTES



सत्यमेव जयते
Government of India

Ministry of Health and Family Welfare

NATIONAL FAMILY HEALTH SURVEY - 5

2019-20

DISTRICT FACT SHEET

EAST KHASI HILLS MEGHALAYA



(स्थापना / Established in 1956)

बेहतर भविष्य के लिए क्षमता निर्माण
Capacity Building for a Better Future

International Institute for Population Sciences
(Deemed University)

Introduction

The National Family Health Survey 2019-20 (NFHS-5), the fifth in the NFHS series, provides information on population, health, and nutrition for India and each state/union territory (UT). Like NFHS-4, NFHS-5 also provides district-level estimates for many important indicators.

The contents of NFHS-5 are similar to NFHS-4 to allow comparisons over time. However, NFHS-5 includes some new topics, such as preschool education, disability, access to a toilet facility, death registration, bathing practices during menstruation, and methods and reasons for abortion. The scope of clinical, anthropometric, and biochemical testing (CAB) has also been expanded to include measurement of waist and hip circumferences, and the age range for the measurement of blood pressure and blood glucose has been expanded. However, HIV testing has been dropped. The NFHS-5 sample has been designed to provide national, state/union territory (UT), and district level estimates of various indicators covered in the survey. However, estimates of indicators of sexual behaviour; husband's background and woman's work; HIV/AIDS knowledge, attitudes and behaviour; and domestic violence are available only at the state/union territory (UT) and national level.

As in the earlier rounds, the Ministry of Health and Family Welfare, Government of India, designated the International Institute for Population Sciences, Mumbai, as the nodal agency to conduct NFHS-5. The main objective of each successive round of the NFHS has been to provide high-quality data on health and family welfare and emerging issues in this area. NFHS-5 data will be useful in setting benchmarks and examining the progress the health sector has made over time. Besides providing evidence for the effectiveness of ongoing programmes, the data from NFHS-5 help in identifying the need for new programmes with an area specific focus and identifying groups that are most in need of essential services.

Four Survey Schedules - Household, Woman's, Man's, and Biomarker - were canvassed in local languages using Computer Assisted Personal Interviewing (CAPI). In the Household Schedule, information was collected on all usual members of the household and visitors who stayed in the household the previous night, as well as socio-economic characteristics of the household; water, sanitation, and hygiene; health insurance coverage; disabilities; land ownership; number of deaths in the household in the three years preceding the survey; and the ownership and use of mosquito nets. The Woman's Schedule covered a wide variety of topics, including the woman's characteristics, marriage, fertility, contraception, children's immunizations and healthcare, nutrition, reproductive health, sexual behaviour, HIV/AIDS, women's empowerment, and domestic violence. The Man's Schedule covered the man's characteristics, marriage, his number of children, contraception, fertility preferences, nutrition, sexual behaviour, health issues, attitudes towards gender roles, and HIV/AIDS. The Biomarker Schedule covered measurements of height, weight, and haemoglobin levels for children; measurements of height, weight, waist and hip circumference, and haemoglobin levels for women age 15-49 years and men age 15-54 years; and blood pressure and random blood glucose levels for women and men age 15 years and over. In addition, women and men were requested to provide a few additional drops of blood from a finger prick for laboratory testing for HbA1c, malaria parasites, and Vitamin D3.

Readers should be cautious while interpreting and comparing the trends as some States/UTs may have smaller sample size. Moreover, at the time of survey, Ayushman Bharat AB-PMJAY and Pradhan Mantri Surakshit Matritva Abhiyan (PMSMA) were not fully rolled out and hence, their coverage may not have been factored in the results of indicator 12 (percentage of households with any usual member covered under a health insurance/financing scheme) and indicator 33 (percentage of mothers who received 4 or more antenatal care check-ups).

This fact sheet provides information on key indicators and trends for East Khasi Hills. NFHS-5 fieldwork for Meghalaya was conducted from 8 July, 2019 to 15 November, 2019 by IQVIA Consulting and Information Services India Pvt. Ltd. In East Khasi Hills, information was gathered from 918 households, 1,066 women, and 131 men.

East Khasi Hills, Meghalaya - Key Indicators

Indicators	NFHS-5 (2019-20)	NFHS-4 (2015-16)
Population and Household Profile		
	Total	Total
1. Female population age 6 years and above who ever attended school (%)	88.7	90.3
2. Population below age 15 years (%)	34.3	34.0
3. Sex ratio of the total population (females per 1,000 males)	1,049	1,027
4. Sex ratio at birth for children born in the last five years (females per 1,000 males)	851	923
5. Children under age 5 years whose birth was registered with the civil authority (%)	77.4	84.9
6. Deaths in the last 3 years registered with the civil authority (%)	55.3	na
7. Population living in households with electricity (%)	96.6	96.0
8. Population living in households with an improved drinking-water source ¹ (%)	90.0	90.4
9. Population living in households that use an improved sanitation facility ² (%)	80.9	67.8
10. Households using clean fuel for cooking ³ (%)	47.4	43.1
11. Households using iodized salt (%)	93.3	99.3
12. Households with any usual member covered under a health insurance/financing scheme (%)	47.1	24.4
13. Children age 5 years who attended pre-primary school during the school year 2019-20 (%)	41.3	na
Characteristics of Women (age 15-49 years)		
14. Women who are literate ⁴ (%)	93.7	na
15. Women with 10 or more years of schooling (%)	43.5	44.0
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	14.6	8.8
17. Births in the 5 years preceding the survey that are third or higher order (%)	4.2	4.0
18. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	8.2	6.0
19. Women age 15-24 years who use hygienic methods of protection during their menstrual period ⁵ (%)	70.7	66.3
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ⁶ (%)	12.1	24.9
21. Any modern method ⁶ (%)	10.6	20.4
22. Female sterilization (%)	4.3	11.8
23. Male sterilization (%)	0.0	0.0
24. IUD/PPIUD (%)	2.4	2.8
25. Pill (%)	1.8	3.1
26. Condom (%)	1.6	2.2
27. Injectables (%)	0.2	0.5
Unmet Need for Family Planning (currently married women age 15-49 years)		
28. Total unmet need ⁷ (%)	33.0	23.5
29. Unmet need for spacing ⁷ (%)	25.2	17.2
Quality of Family Planning Services		
30. Health worker ever talked to female non-users about family planning (%)	22.7	22.6
31. Current users ever told about side effects of current method ⁸ (%)	*	69.3

Note: Major indicators are highlighted in grey.

LHV = Lady health visitor, ANM = Auxiliary nurse midwife

na = Not available

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rainwater, tanker truck, cart with small tank, bottled water, community RO plant.

²Flush to piped sewer system, flush to septic tank, flush to pit latrine, flush to don't know where, ventilated improved pit (VIP)/biogas latrine, pit latrine with slab, twin pit/composting toilet, which is not shared with any other household. This indicator does not denote access to toilet facility completely.

³Electricity, LPG/natural gas, biogas.

⁴Refers to women who completed standard 9 or higher and women who can read a whole sentence or part of a sentence.

⁵Locally prepared napkins, sanitary napkins, tampons, and menstrual cups are considered to be hygienic methods of protection.

⁶Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

⁷Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.
- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

Women who are classified as infecund have no unmet need because they are not at risk of becoming pregnant. Unmet need for family planning is the sum of unmet need for spacing plus unmet need for limiting.

⁸Based on current users of female sterilization, IUD/PPIUD, injectables, and pills who started using that method in the past 5 years.

East Khasi Hills, Meghalaya - Key Indicators

Indicators	NFHS-5 (2019-20)	NFHS-4 (2015-16)
Maternal and Child Health	Total	Total
Maternity Care (for last birth in the 5 years before the survey)		
32. Mothers who had an antenatal check-up in the first trimester (%)	56.7	72.5
33. Mothers who had at least 4 antenatal care visits (%)	57.5	72.7
34. Mothers whose last birth was protected against neonatal tetanus ⁹ (%)	80.2	84.2
35. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	40.1	56.4
36. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	27.1	24.3
37. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	87.8	88.6
38. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery (%)	33.6	64.8
39. Average out-of-pocket expenditure per delivery in a public health facility (Rs.)	3,258	2,519
40. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	3.7	4.3
41. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery (%)	41.6	na
Delivery Care (for births in the 5 years before the survey)		
42. Institutional births (%)	63.4	66.8
43. Institutional births in public facility (%)	48.4	41.3
44. Home births that were conducted by skilled health personnel ¹⁰ (%)	6.2	2.6
45. Births attended by skilled health personnel ¹⁰ (%)	68.7	69.5
46. Births delivered by caesarean section (%)	13.9	14.2
47. Births in a private health facility that were delivered by caesarean section (%)	48.3	30.0
48. Births in a public health facility that were delivered by caesarean section (%)	13.7	16.0
Child Vaccinations and Vitamin A Supplementation		
49. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ¹¹ (%)	49.1	68.7
50. Children age 12-23 months fully vaccinated based on information from vaccination card only ¹² (%)	69.4	87.4
51. Children age 12-23 months who have received BCG (%)	84.2	92.2
52. Children age 12-23 months who have received 3 doses of polio vaccine ¹³ (%)	53.2	75.3
53. Children age 12-23 months who have received 3 doses of penta or DPT vaccine (%)	59.2	76.4
54. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	56.6	78.8
55. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	10.9	na
56. Children age 12-23 months who have received 3 doses of rotavirus vaccine ¹⁴ (%)	5.2	na
57. Children age 12-23 months who have received 3 doses of penta or hepatitis B vaccine (%)	46.8	67.3
58. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	54.4	64.5
59. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	87.5	83.3
60. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	9.9	12.3
Treatment of Childhood Diseases (children under age 5 years)		
61. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	11.5	8.7
62. Children with diarrhoea in the 2 weeks preceding the survey who received oral rehydration salts (ORS) (%)	(74.3)	82.0
63. Children with diarrhoea in the 2 weeks preceding the survey who received zinc (%)	(31.2)	52.9
64. Children with diarrhoea in the 2 weeks preceding the survey taken to a health facility or health provider (%)	(71.3)	81.8
65. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	5.1	3.1
66. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	72.7	87.7

⁹Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

¹⁰Doctor/nurse/LHV/ANM/midwife/other health personnel.

¹¹Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

¹²Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

¹³Not including polio vaccination given at birth.

¹⁴Since rotavirus is not being provided across all states and districts, the levels should not be compared.

East Khasi Hills, Meghalaya - Key Indicators

Indicators	NFHS-5 (2019-20)	NFHS-4 (2015-16)
Child Feeding Practices and Nutritional Status of Children		
67. Children under age 3 years breastfed within one hour of birth ¹⁵ (%)	86.3	52.0
68. Children under age 6 months exclusively breastfed ¹⁶ (%)	(25.1)	34.5
69. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁶ (%)	*	(60.1)
70. Breastfeeding children age 6-23 months receiving an adequate diet ^{16, 17} (%)	32.5	28.8
71. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{16, 17} (%)	(41.3)	26.2
72. Total children age 6-23 months receiving an adequate diet ^{16, 17} (%)	34.8	28.2
73. Children under 5 years who are stunted (height-for-age) ¹⁸ (%)	44.6	48.6
74. Children under 5 years who are wasted (weight-for-height) ¹⁸ (%)	11.2	11.1
75. Children under 5 years who are severely wasted (weight-for-height) ¹⁹ (%)	4.9	3.7
76. Children under 5 years who are underweight (weight-for-age) ¹⁸ (%)	23.8	30.4
77. Children under 5 years who are overweight (weight-for-height) ²⁰ (%)	4.5	2.9
Nutritional Status of Women (age 15-49 years)		
78. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ²¹ (%)	11.6	12.8
79. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ²¹ (%)	15.6	16.7
80. Women who have high risk waist-to-hip ratio (≥0.85) (%)	49.3	na
Anaemia among Children and Women		
81. Children age 6-59 months who are anaemic (<11.0 g/dl) ²² (%)	39.0	46.5
82. Non-pregnant women age 15-49 years who are anaemic (<12.0 g/dl) ²² (%)	48.5	46.7
83. Pregnant women age 15-49 years who are anaemic (<11.0 g/dl) ²² (%)	43.7	47.7
84. All women age 15-49 years who are anaemic ²² (%)	48.2	46.8
85. All women age 15-19 years who are anaemic ²² (%)	40.0	42.5
Blood Sugar Level among Adults (age 15 years and above)		
Women		
86. Blood sugar level - high (141-160 mg/dl) ²³ (%)	2.7	na
87. Blood sugar level - very high (>160 mg/dl) ²³ (%)	3.5	na
88. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ²³ (%)	9.1	na
Men		
89. Blood sugar level - high (141-160 mg/dl) ²³ (%)	3.6	na
90. Blood sugar level - very high (>160 mg/dl) ²³ (%)	3.4	na
91. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ²³ (%)	9.4	na
Hypertension among Adults (age 15 years and above)		
Women		
92. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	11.2	na
93. Moderately or severely elevated blood pressure (Systolic ≥160mm of Hg and/or Diastolic ≥100mm of Hg) (%)	5.6	na
94. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	24.5	na
Men		
95. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	14.5	na
96. Moderately or severely elevated blood pressure (Systolic ≥160mm of Hg and/or Diastolic ≥100mm of Hg) (%)	4.3	na
97. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	25.0	na
Screening for Cancer among Women (age 30-49 years)		
98. Ever undergone a screening test for cervical cancer (%)	1.3	na
99. Ever undergone a breast examination for breast cancer (%)	1.3	na
100. Ever undergone an oral cavity examination for oral cancer (%)	0.8	na
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
101. Women age 15 years and above who use any kind of tobacco (%)	29.8	na
102. Men age 15 years and above who use any kind of tobacco (%)	51.9	na
103. Women age 15 years and above who consume alcohol (%)	1.2	na
104. Men age 15 years and above who consume alcohol (%)	24.0	na

¹⁵Based on the last child born in the 3 years before the survey.

¹⁶Based on the youngest child living with the mother.

¹⁷Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹⁸Below -2 standard deviations, based on the WHO standard.

¹⁹Below -3 standard deviations, based on the WHO standard.

²⁰Above +2 standard deviations, based on the WHO standard.

²¹Excludes pregnant women and women with a birth in the preceding 2 months.

²²Haemoglobin in grams per decilitre (g/dl). Among children, prevalence is adjusted for altitude. Among women, prevalence is adjusted for altitude and for smoking status, if known. As NFHS uses the capillary blood for estimation of anaemia, the results of NFHS-5 need not be compared with other surveys using venous blood.

²³Random blood sugar measurement.

NOTES



सत्यमेव जयते
Government of India

Ministry of Health and Family Welfare

NATIONAL FAMILY HEALTH SURVEY - 5

2019-20

DISTRICT FACT SHEET

NORTH GARO HILLS

MEGHALAYA



(स्थापना / Established in 1956)

बेहतर भविष्य के लिए क्षमता निर्माण
Capacity Building for a Better Future

International Institute for Population Sciences
(Deemed University)

Introduction

The National Family Health Survey 2019-20 (NFHS-5), the fifth in the NFHS series, provides information on population, health, and nutrition for India and each state/union territory (UT). Like NFHS-4, NFHS-5 also provides district-level estimates for many important indicators.

The contents of NFHS-5 are similar to NFHS-4 to allow comparisons over time. However, NFHS-5 includes some new topics, such as preschool education, disability, access to a toilet facility, death registration, bathing practices during menstruation, and methods and reasons for abortion. The scope of clinical, anthropometric, and biochemical testing (CAB) has also been expanded to include measurement of waist and hip circumferences, and the age range for the measurement of blood pressure and blood glucose has been expanded. However, HIV testing has been dropped. The NFHS-5 sample has been designed to provide national, state/union territory (UT), and district level estimates of various indicators covered in the survey. However, estimates of indicators of sexual behaviour; husband's background and woman's work; HIV/AIDS knowledge, attitudes and behaviour; and domestic violence are available only at the state/union territory (UT) and national level.

As in the earlier rounds, the Ministry of Health and Family Welfare, Government of India, designated the International Institute for Population Sciences, Mumbai, as the nodal agency to conduct NFHS-5. The main objective of each successive round of the NFHS has been to provide high-quality data on health and family welfare and emerging issues in this area. NFHS-5 data will be useful in setting benchmarks and examining the progress the health sector has made over time. Besides providing evidence for the effectiveness of ongoing programmes, the data from NFHS-5 help in identifying the need for new programmes with an area specific focus and identifying groups that are most in need of essential services.

Four Survey Schedules - Household, Woman's, Man's, and Biomarker - were canvassed in local languages using Computer Assisted Personal Interviewing (CAPI). In the Household Schedule, information was collected on all usual members of the household and visitors who stayed in the household the previous night, as well as socio-economic characteristics of the household; water, sanitation, and hygiene; health insurance coverage; disabilities; land ownership; number of deaths in the household in the three years preceding the survey; and the ownership and use of mosquito nets. The Woman's Schedule covered a wide variety of topics, including the woman's characteristics, marriage, fertility, contraception, children's immunizations and healthcare, nutrition, reproductive health, sexual behaviour, HIV/AIDS, women's empowerment, and domestic violence. The Man's Schedule covered the man's characteristics, marriage, his number of children, contraception, fertility preferences, nutrition, sexual behaviour, health issues, attitudes towards gender roles, and HIV/AIDS. The Biomarker Schedule covered measurements of height, weight, and haemoglobin levels for children; measurements of height, weight, waist and hip circumference, and haemoglobin levels for women age 15-49 years and men age 15-54 years; and blood pressure and random blood glucose levels for women and men age 15 years and over. In addition, women and men were requested to provide a few additional drops of blood from a finger prick for laboratory testing for HbA1c, malaria parasites, and Vitamin D3.

Readers should be cautious while interpreting and comparing the trends as some States/UTs may have smaller sample size. Moreover, at the time of survey, Ayushman Bharat AB-PMJAY and Pradhan Mantri Surakshit Matritva Abhiyan (PMSMA) were not fully rolled out and hence, their coverage may not have been factored in the results of indicator 12 (percentage of households with any usual member covered under a health insurance/financing scheme) and indicator 33 (percentage of mothers who received 4 or more antenatal care check-ups).

This fact sheet provides information on key indicators for North Garo Hills. NFHS-5 fieldwork for Meghalaya was conducted from 8 July, 2019 to 15 November, 2019 by IQVIA Consulting and Information Services India Pvt. Ltd. In North Garo Hills, information was gathered from 923 households, 1,283 women, and 180 men.

North Garo Hills, Meghalaya - Key Indicators

Indicators	NFHS-5 (2019-20)
Population and Household Profile	
	Total
1. Female population age 6 years and above who ever attended school (%)	91.3
2. Population below age 15 years (%)	29.2
3. Sex ratio of the total population (females per 1,000 males)	1,009
4. Sex ratio at birth for children born in the last five years (females per 1,000 males)	894
5. Children under age 5 years whose birth was registered with the civil authority (%)	79.5
6. Deaths in the last 3 years registered with the civil authority (%)	(43.8)
7. Population living in households with electricity (%)	97.5
8. Population living in households with an improved drinking-water source ¹ (%)	51.5
9. Population living in households that use an improved sanitation facility ² (%)	84.9
10. Households using clean fuel for cooking ³ (%)	24.4
11. Households using iodized salt (%)	84.1
12. Households with any usual member covered under a health insurance/financing scheme (%)	60.8
13. Children age 5 years who attended pre-primary school during the school year 2019-20 (%)	12.7
Characteristics of Women (age 15-49 years)	
14. Women who are literate ⁴ (%)	86.1
15. Women with 10 or more years of schooling (%)	38.4
Marriage and Fertility	
16. Women age 20-24 years married before age 18 years (%)	13.2
17. Births in the 5 years preceding the survey that are third or higher order (%)	1.1
18. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	3.7
19. Women age 15-24 years who use hygienic methods of protection during their menstrual period ⁵ (%)	74.9
Current Use of Family Planning Methods (currently married women age 15-49 years)	
20. Any method ⁶ (%)	19.2
21. Any modern method ⁶ (%)	17.7
22. Female sterilization (%)	2.9
23. Male sterilization (%)	0.0
24. IUD/PPIUD (%)	3.4
25. Pill (%)	7.4
26. Condom (%)	3.2
27. Injectables (%)	0.7
Unmet Need for Family Planning (currently married women age 15-49 years)	
28. Total unmet need ⁷ (%)	25.2
29. Unmet need for spacing ⁷ (%)	12.9
Quality of Family Planning Services	
30. Health worker ever talked to female non-users about family planning (%)	8.0
31. Current users ever told about side effects of current method ⁸ (%)	70.4

Note: Indicator estimates for NFHS-4 are not shown in this table since no comparable estimates are available from NFHS-4 in this district due to district boundary changes or a newly formed district. Major indicators are highlighted in grey.

LHV = Lady health visitor, ANM = Auxiliary nurse midwife

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rainwater, tanker truck, cart with small tank, bottled water, community RO plant.

²Flush to piped sewer system, flush to septic tank, flush to pit latrine, flush to don't know where, ventilated improved pit (VIP)/biogas latrine, pit latrine with slab, twin pit/composting toilet, which is not shared with any other household. This indicator does not denote access to toilet facility completely.

³Electricity, LPG/natural gas, biogas.

⁴Refers to women who completed standard 9 or higher and women who can read a whole sentence or part of a sentence.

⁵Locally prepared napkins, sanitary napkins, tampons, and menstrual cups are considered to be hygienic methods of protection.

⁶Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

⁷Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.
- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

Women who are classified as infecund have no unmet need because they are not at risk of becoming pregnant. Unmet need for family planning is the sum of unmet need for spacing plus unmet need for limiting.

⁸Based on current users of female sterilization, IUD/PPIUD, injectables, and pills who started using that method in the past 5 years.

North Garo Hills, Meghalaya - Key Indicators

Indicators	NFHS-5 (2019-20)
Maternal and Child Health	Total
Maternity Care (for last birth in the 5 years before the survey)	
32. Mothers who had an antenatal check-up in the first trimester (%)	40.3
33. Mothers who had at least 4 antenatal care visits (%)	24.3
34. Mothers whose last birth was protected against neonatal tetanus ⁹ (%)	55.5
35. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	23.7
36. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	7.9
37. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	85.8
38. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery (%)	40.5
39. Average out-of-pocket expenditure per delivery in a public health facility (Rs.)	4,504
40. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	1.0
41. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery (%)	41.8
Delivery Care (for births in the 5 years before the survey)	
42. Institutional births (%)	61.5
43. Institutional births in public facility (%)	60.4
44. Home births that were conducted by skilled health personnel ¹⁰ (%)	5.9
45. Births attended by skilled health personnel ¹⁰ (%)	63.2
46. Births delivered by caesarean section (%)	3.3
47. Births in a private health facility that were delivered by caesarean section (%)	*
48. Births in a public health facility that were delivered by caesarean section (%)	5.4
Child Vaccinations and Vitamin A Supplementation	
49. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ¹¹ (%)	(47.5)
50. Children age 12-23 months fully vaccinated based on information from vaccination card only ¹² (%)	*
51. Children age 12-23 months who have received BCG (%)	(65.9)
52. Children age 12-23 months who have received 3 doses of polio vaccine ¹³ (%)	(54.7)
53. Children age 12-23 months who have received 3 doses of penta or DPT vaccine (%)	(57.9)
54. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	(58.1)
55. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	(2.8)
56. Children age 12-23 months who have received 3 doses of rotavirus vaccine ¹⁴ (%)	(8.2)
57. Children age 12-23 months who have received 3 doses of penta or hepatitis B vaccine (%)	(39.5)
58. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	42.6
59. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	(96.9)
60. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	(0.0)
Treatment of Childhood Diseases (children under age 5 years)	
61. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	6.6
62. Children with diarrhoea in the 2 weeks preceding the survey who received oral rehydration salts (ORS) (%)	*
63. Children with diarrhoea in the 2 weeks preceding the survey who received zinc (%)	*
64. Children with diarrhoea in the 2 weeks preceding the survey taken to a health facility or health provider (%)	*
65. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	0.9
66. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	56.6

⁹Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

¹⁰Doctor/nurse/LHV/ANM/midwife/other health personnel.

¹¹Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

¹²Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

¹³Not including polio vaccination given at birth.

¹⁴Since rotavirus is not being provided across all states and districts, the levels should not be compared.

North Garo Hills, Meghalaya - Key Indicators

Indicators	NFHS-5 (2019-20)
Child Feeding Practices and Nutritional Status of Children	
Total	
67. Children under age 3 years breastfed within one hour of birth ¹⁵ (%)	62.3
68. Children under age 6 months exclusively breastfed ¹⁶ (%)	*
69. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁶ (%)	*
70. Breastfeeding children age 6-23 months receiving an adequate diet ^{16, 17} (%)	11.7
71. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{16, 17} (%)	*
72. Total children age 6-23 months receiving an adequate diet ^{16, 17} (%)	13.2
73. Children under 5 years who are stunted (height-for-age) ¹⁸ (%)	35.6
74. Children under 5 years who are wasted (weight-for-height) ¹⁸ (%)	11.7
75. Children under 5 years who are severely wasted (weight-for-height) ¹⁹ (%)	5.6
76. Children under 5 years who are underweight (weight-for-age) ¹⁸ (%)	15.8
77. Children under 5 years who are overweight (weight-for-height) ²⁰ (%)	7.3
Nutritional Status of Women (age 15-49 years)	
78. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ²¹ (%)	7.7
79. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ²¹ (%)	11.1
80. Women who have high risk waist-to-hip ratio (≥0.85) (%)	65.9
Anaemia among Children and Women	
81. Children age 6-59 months who are anaemic (<11.0 g/dl) ²² (%)	30.2
82. Non-pregnant women age 15-49 years who are anaemic (<12.0 g/dl) ²² (%)	58.9
83. Pregnant women age 15-49 years who are anaemic (<11.0 g/dl) ²² (%)	(40.8)
84. All women age 15-49 years who are anaemic ²² (%)	58.4
85. All women age 15-19 years who are anaemic ²² (%)	65.2
Blood Sugar Level among Adults (age 15 years and above)	
Women	
86. Blood sugar level - high (141-160 mg/dl) ²³ (%)	7.8
87. Blood sugar level - very high (>160 mg/dl) ²³ (%)	2.7
88. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ²³ (%)	12.2
Men	
89. Blood sugar level - high (141-160 mg/dl) ²³ (%)	11.8
90. Blood sugar level - very high (>160 mg/dl) ²³ (%)	4.1
91. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ²³ (%)	18.7
Hypertension among Adults (age 15 years and above)	
Women	
92. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	10.5
93. Moderately or severely elevated blood pressure (Systolic ≥160mm of Hg and/or Diastolic ≥100mm of Hg) (%)	4.6
94. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	20.0
Men	
95. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	15.7
96. Moderately or severely elevated blood pressure (Systolic ≥160mm of Hg and/or Diastolic ≥100mm of Hg) (%)	5.3
97. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	25.4
Screening for Cancer among Women (age 30-49 years)	
98. Ever undergone a screening test for cervical cancer (%)	0.0
99. Ever undergone a breast examination for breast cancer (%)	0.0
100. Ever undergone an oral cavity examination for oral cancer (%)	0.2
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)	
101. Women age 15 years and above who use any kind of tobacco (%)	3.8
102. Men age 15 years and above who use any kind of tobacco (%)	54.1
103. Women age 15 years and above who consume alcohol (%)	2.2
104. Men age 15 years and above who consume alcohol (%)	40.2

¹⁵Based on the last child born in the 3 years before the survey.

¹⁶Based on the youngest child living with the mother.

¹⁷Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹⁸Below -2 standard deviations, based on the WHO standard.

¹⁹Below -3 standard deviations, based on the WHO standard.

²⁰Above +2 standard deviations, based on the WHO standard.

²¹Excludes pregnant women and women with a birth in the preceding 2 months.

²²Haemoglobin in grams per decilitre (g/dl). Among children, prevalence is adjusted for altitude. Among women, prevalence is adjusted for altitude and for smoking status, if known. As NFHS uses the capillary blood for estimation of anaemia, the results of NFHS-5 need not be compared with other surveys using venous blood.

²³Random blood sugar measurement.

NOTES



सत्यमेव जयते
Government of India

Ministry of Health and Family Welfare

NATIONAL FAMILY HEALTH SURVEY - 5

2019-20

DISTRICT FACT SHEET

RIBHOI
MEGHALAYA



(स्थापना / Established in 1956)

बेहतर भविष्य के लिए क्षमता निर्माण
Capacity Building for a Better Future

International Institute for Population Sciences
(Deemed University)

Introduction

The National Family Health Survey 2019-20 (NFHS-5), the fifth in the NFHS series, provides information on population, health, and nutrition for India and each state/union territory (UT). Like NFHS-4, NFHS-5 also provides district-level estimates for many important indicators.

The contents of NFHS-5 are similar to NFHS-4 to allow comparisons over time. However, NFHS-5 includes some new topics, such as preschool education, disability, access to a toilet facility, death registration, bathing practices during menstruation, and methods and reasons for abortion. The scope of clinical, anthropometric, and biochemical testing (CAB) has also been expanded to include measurement of waist and hip circumferences, and the age range for the measurement of blood pressure and blood glucose has been expanded. However, HIV testing has been dropped. The NFHS-5 sample has been designed to provide national, state/union territory (UT), and district level estimates of various indicators covered in the survey. However, estimates of indicators of sexual behaviour; husband's background and woman's work; HIV/AIDS knowledge, attitudes and behaviour; and domestic violence are available only at the state/union territory (UT) and national level.

As in the earlier rounds, the Ministry of Health and Family Welfare, Government of India, designated the International Institute for Population Sciences, Mumbai, as the nodal agency to conduct NFHS-5. The main objective of each successive round of the NFHS has been to provide high-quality data on health and family welfare and emerging issues in this area. NFHS-5 data will be useful in setting benchmarks and examining the progress the health sector has made over time. Besides providing evidence for the effectiveness of ongoing programmes, the data from NFHS-5 help in identifying the need for new programmes with an area specific focus and identifying groups that are most in need of essential services.

Four Survey Schedules - Household, Woman's, Man's, and Biomarker - were canvassed in local languages using Computer Assisted Personal Interviewing (CAPI). In the Household Schedule, information was collected on all usual members of the household and visitors who stayed in the household the previous night, as well as socio-economic characteristics of the household; water, sanitation, and hygiene; health insurance coverage; disabilities; land ownership; number of deaths in the household in the three years preceding the survey; and the ownership and use of mosquito nets. The Woman's Schedule covered a wide variety of topics, including the woman's characteristics, marriage, fertility, contraception, children's immunizations and healthcare, nutrition, reproductive health, sexual behaviour, HIV/AIDS, women's empowerment, and domestic violence. The Man's Schedule covered the man's characteristics, marriage, his number of children, contraception, fertility preferences, nutrition, sexual behaviour, health issues, attitudes towards gender roles, and HIV/AIDS. The Biomarker Schedule covered measurements of height, weight, and haemoglobin levels for children; measurements of height, weight, waist and hip circumference, and haemoglobin levels for women age 15-49 years and men age 15-54 years; and blood pressure and random blood glucose levels for women and men age 15 years and over. In addition, women and men were requested to provide a few additional drops of blood from a finger prick for laboratory testing for HbA1c, malaria parasites, and Vitamin D3.

Readers should be cautious while interpreting and comparing the trends as some States/UTs may have smaller sample size. Moreover, at the time of survey, Ayushman Bharat AB-PMJAY and Pradhan Mantri Surakshit Matritva Abhiyan (PMSMA) were not fully rolled out and hence, their coverage may not have been factored in the results of indicator 12 (percentage of households with any usual member covered under a health insurance/financing scheme) and indicator 33 (percentage of mothers who received 4 or more antenatal care check-ups).

This fact sheet provides information on key indicators and trends for Ribhoi. NFHS-5 fieldwork for Meghalaya was conducted from 8 July, 2019 to 15 November, 2019 by IQVIA Consulting and Information Services India Pvt. Ltd. In Ribhoi, information was gathered from 927 households, 1,238 women, and 145 men.

Ribhoi, Meghalaya - Key Indicators

Indicators	NFHS-5 (2019-20)	NFHS-4 (2015-16)
Population and Household Profile		
	Total	Total
1. Female population age 6 years and above who ever attended school (%)	85.3	79.3
2. Population below age 15 years (%)	40.4	40.0
3. Sex ratio of the total population (females per 1,000 males)	998	1,007
4. Sex ratio at birth for children born in the last five years (females per 1,000 males)	871	1,113
5. Children under age 5 years whose birth was registered with the civil authority (%)	85.9	83.0
6. Deaths in the last 3 years registered with the civil authority (%)	51.1	na
7. Population living in households with electricity (%)	92.1	86.7
8. Population living in households with an improved drinking-water source ¹ (%)	86.2	70.5
9. Population living in households that use an improved sanitation facility ² (%)	85.3	58.1
10. Households using clean fuel for cooking ³ (%)	21.0	9.6
11. Households using iodized salt (%)	96.6	97.8
12. Households with any usual member covered under a health insurance/financing scheme (%)	66.8	36.3
13. Children age 5 years who attended pre-primary school during the school year 2019-20 (%)	28.7	na
Characteristics of Women (age 15-49 years)		
14. Women who are literate ⁴ (%)	89.5	na
15. Women with 10 or more years of schooling (%)	23.1	14.9
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	20.3	20.6
17. Births in the 5 years preceding the survey that are third or higher order (%)	5.5	5.8
18. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	6.8	10.9
19. Women age 15-24 years who use hygienic methods of protection during their menstrual period ⁵ (%)	67.8	56.3
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ⁶ (%)	33.1	23.7
21. Any modern method ⁶ (%)	19.7	22.8
22. Female sterilization (%)	6.3	6.6
23. Male sterilization (%)	0.0	0.0
24. IUD/PPIUD (%)	4.9	2.0
25. Pill (%)	3.7	11.1
26. Condom (%)	4.4	2.5
27. Injectables (%)	0.2	0.4
Unmet Need for Family Planning (currently married women age 15-49 years)		
28. Total unmet need ⁷ (%)	27.0	23.9
29. Unmet need for spacing ⁷ (%)	18.4	18.2
Quality of Family Planning Services		
30. Health worker ever talked to female non-users about family planning (%)	34.2	20.3
31. Current users ever told about side effects of current method ⁸ (%)	71.1	43.3

Note: Major indicators are highlighted in grey.

LHV = Lady health visitor, ANM = Auxiliary nurse midwife

na = Not available

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rainwater, tanker truck, cart with small tank, bottled water, community RO plant.

²Flush to piped sewer system, flush to septic tank, flush to pit latrine, flush to don't know where, ventilated improved pit (VIP)/biogas latrine, pit latrine with slab, twin pit/composting toilet, which is not shared with any other household. This indicator does not denote access to toilet facility completely.

³Electricity, LPG/natural gas, biogas.

⁴Refers to women who completed standard 9 or higher and women who can read a whole sentence or part of a sentence.

⁵Locally prepared napkins, sanitary napkins, tampons, and menstrual cups are considered to be hygienic methods of protection.

⁶Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

⁷Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.
- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

Women who are classified as infecund have no unmet need because they are not at risk of becoming pregnant. Unmet need for family planning is the sum of unmet need for spacing plus unmet need for limiting.

⁸Based on current users of female sterilization, IUD/PPIUD, injectables, and pills who started using that method in the past 5 years.

Ribhoi, Meghalaya - Key Indicators

Indicators	NFHS-5 (2019-20)	NFHS-4 (2015-16)
Maternal and Child Health	Total	Total
Maternity Care (for last birth in the 5 years before the survey)		
32. Mothers who had an antenatal check-up in the first trimester (%)	56.1	43.8
33. Mothers who had at least 4 antenatal care visits (%)	61.0	40.5
34. Mothers whose last birth was protected against neonatal tetanus ⁹ (%)	86.2	75.8
35. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	48.8	33.5
36. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	22.4	11.8
37. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	97.5	97.2
38. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery (%)	39.5	45.0
39. Average out-of-pocket expenditure per delivery in a public health facility (Rs.)	3,648	2,206
40. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	3.0	1.2
41. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery (%)	39.3	na
Delivery Care (for births in the 5 years before the survey)		
42. Institutional births (%)	56.9	44.9
43. Institutional births in public facility (%)	45.6	38.3
44. Home births that were conducted by skilled health personnel ¹⁰ (%)	8.4	2.2
45. Births attended by skilled health personnel ¹⁰ (%)	65.8	46.4
46. Births delivered by caesarean section (%)	8.7	5.8
47. Births in a private health facility that were delivered by caesarean section (%)	36.6	(43.5)
48. Births in a public health facility that were delivered by caesarean section (%)	10.0	7.7
Child Vaccinations and Vitamin A Supplementation		
49. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ¹¹ (%)	74.8	55.7
50. Children age 12-23 months fully vaccinated based on information from vaccination card only ¹² (%)	86.7	80.5
51. Children age 12-23 months who have received BCG (%)	94.0	84.4
52. Children age 12-23 months who have received 3 doses of polio vaccine ¹³ (%)	77.1	63.8
53. Children age 12-23 months who have received 3 doses of penta or DPT vaccine (%)	82.8	71.5
54. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	80.9	73.2
55. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	11.8	na
56. Children age 12-23 months who have received 3 doses of rotavirus vaccine ¹⁴ (%)	4.4	na
57. Children age 12-23 months who have received 3 doses of penta or hepatitis B vaccine (%)	71.5	55.6
58. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	66.6	66.5
59. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	92.6	100.0
60. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	4.1	0.0
Treatment of Childhood Diseases (children under age 5 years)		
61. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	11.3	14.7
62. Children with diarrhoea in the 2 weeks preceding the survey who received oral rehydration salts (ORS) (%)	69.6	82.4
63. Children with diarrhoea in the 2 weeks preceding the survey who received zinc (%)	42.1	53.1
64. Children with diarrhoea in the 2 weeks preceding the survey taken to a health facility or health provider (%)	53.5	79.3
65. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	3.6	5.0
66. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	68.7	81.1

⁹Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

¹⁰Doctor/nurse/LHV/ANM/midwife/other health personnel.

¹¹Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

¹²Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

¹³Not including polio vaccination given at birth.

¹⁴Since rotavirus is not being provided across all states and districts, the levels should not be compared.

Ribhoi, Meghalaya - Key Indicators

Indicators	NFHS-5 (2019-20)	NFHS-4 (2015-16)
Child Feeding Practices and Nutritional Status of Children		
67. Children under age 3 years breastfed within one hour of birth ¹⁵ (%)	85.9	76.1
68. Children under age 6 months exclusively breastfed ¹⁶ (%)	36.8	34.1
69. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁶ (%)	(63.5)	(61.4)
70. Breastfeeding children age 6-23 months receiving an adequate diet ^{16, 17} (%)	38.2	33.4
71. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{16, 17} (%)	(34.7)	(8.8)
72. Total children age 6-23 months receiving an adequate diet ^{16, 17} (%)	37.3	28.9
73. Children under 5 years who are stunted (height-for-age) ¹⁸ (%)	42.5	51.6
74. Children under 5 years who are wasted (weight-for-height) ¹⁸ (%)	17.9	9.4
75. Children under 5 years who are severely wasted (weight-for-height) ¹⁹ (%)	7.5	5.1
76. Children under 5 years who are underweight (weight-for-age) ¹⁸ (%)	29.6	30.8
77. Children under 5 years who are overweight (weight-for-height) ²⁰ (%)	4.8	4.9
Nutritional Status of Women (age 15-49 years)		
78. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ²¹ (%)	15.5	16.7
79. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ²¹ (%)	12.2	10.5
80. Women who have high risk waist-to-hip ratio (≥0.85) (%)	59.4	na
Anaemia among Children and Women		
81. Children age 6-59 months who are anaemic (<11.0 g/dl) ²² (%)	45.7	44.4
82. Non-pregnant women age 15-49 years who are anaemic (<12.0 g/dl) ²² (%)	62.2	68.0
83. Pregnant women age 15-49 years who are anaemic (<11.0 g/dl) ²² (%)	65.0	66.7
84. All women age 15-49 years who are anaemic ²² (%)	62.4	68.0
85. All women age 15-19 years who are anaemic ²² (%)	57.1	61.0
Blood Sugar Level among Adults (age 15 years and above)		
Women		
86. Blood sugar level - high (141-160 mg/dl) ²³ (%)	4.6	na
87. Blood sugar level - very high (>160 mg/dl) ²³ (%)	2.8	na
88. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ²³ (%)	8.0	na
Men		
89. Blood sugar level - high (141-160 mg/dl) ²³ (%)	4.9	na
90. Blood sugar level - very high (>160 mg/dl) ²³ (%)	2.7	na
91. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ²³ (%)	8.1	na
Hypertension among Adults (age 15 years and above)		
Women		
92. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	7.4	na
93. Moderately or severely elevated blood pressure (Systolic ≥160mm of Hg and/or Diastolic ≥100mm of Hg) (%)	2.0	na
94. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	13.5	na
Men		
95. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	12.3	na
96. Moderately or severely elevated blood pressure (Systolic ≥160mm of Hg and/or Diastolic ≥100mm of Hg) (%)	2.9	na
97. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	16.0	na
Screening for Cancer among Women (age 30-49 years)		
98. Ever undergone a screening test for cervical cancer (%)	0.6	na
99. Ever undergone a breast examination for breast cancer (%)	0.4	na
100. Ever undergone an oral cavity examination for oral cancer (%)	0.4	na
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
101. Women age 15 years and above who use any kind of tobacco (%)	34.1	na
102. Men age 15 years and above who use any kind of tobacco (%)	60.4	na
103. Women age 15 years and above who consume alcohol (%)	0.7	na
104. Men age 15 years and above who consume alcohol (%)	30.9	na

¹⁵Based on the last child born in the 3 years before the survey.

¹⁶Based on the youngest child living with the mother.

¹⁷Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹⁸Below -2 standard deviations, based on the WHO standard.

¹⁹Below -3 standard deviations, based on the WHO standard.

²⁰Above +2 standard deviations, based on the WHO standard.

²¹Excludes pregnant women and women with a birth in the preceding 2 months.

²²Haemoglobin in grams per decilitre (g/dl). Among children, prevalence is adjusted for altitude. Among women, prevalence is adjusted for altitude and for smoking status, if known. As NFHS uses the capillary blood for estimation of anaemia, the results of NFHS-5 need not be compared with other surveys using venous blood.

²³Random blood sugar measurement.

NOTES



सत्यमेव जयते
Government of India

Ministry of Health and Family Welfare

NATIONAL FAMILY HEALTH SURVEY - 5

2019-20

DISTRICT FACT SHEET

SOUTH GARO HILLS

MEGHALAYA



(स्थापना / Established in 1956)

बेहतर भविष्य के लिए क्षमता निर्माण
Capacity Building for a Better Future

International Institute for Population Sciences
(Deemed University)

Introduction

The National Family Health Survey 2019-20 (NFHS-5), the fifth in the NFHS series, provides information on population, health, and nutrition for India and each state/union territory (UT). Like NFHS-4, NFHS-5 also provides district-level estimates for many important indicators.

The contents of NFHS-5 are similar to NFHS-4 to allow comparisons over time. However, NFHS-5 includes some new topics, such as preschool education, disability, access to a toilet facility, death registration, bathing practices during menstruation, and methods and reasons for abortion. The scope of clinical, anthropometric, and biochemical testing (CAB) has also been expanded to include measurement of waist and hip circumferences, and the age range for the measurement of blood pressure and blood glucose has been expanded. However, HIV testing has been dropped. The NFHS-5 sample has been designed to provide national, state/union territory (UT), and district level estimates of various indicators covered in the survey. However, estimates of indicators of sexual behaviour; husband's background and woman's work; HIV/AIDS knowledge, attitudes and behaviour; and domestic violence are available only at the state/union territory (UT) and national level.

As in the earlier rounds, the Ministry of Health and Family Welfare, Government of India, designated the International Institute for Population Sciences, Mumbai, as the nodal agency to conduct NFHS-5. The main objective of each successive round of the NFHS has been to provide high-quality data on health and family welfare and emerging issues in this area. NFHS-5 data will be useful in setting benchmarks and examining the progress the health sector has made over time. Besides providing evidence for the effectiveness of ongoing programmes, the data from NFHS-5 help in identifying the need for new programmes with an area specific focus and identifying groups that are most in need of essential services.

Four Survey Schedules - Household, Woman's, Man's, and Biomarker - were canvassed in local languages using Computer Assisted Personal Interviewing (CAPI). In the Household Schedule, information was collected on all usual members of the household and visitors who stayed in the household the previous night, as well as socio-economic characteristics of the household; water, sanitation, and hygiene; health insurance coverage; disabilities; land ownership; number of deaths in the household in the three years preceding the survey; and the ownership and use of mosquito nets. The Woman's Schedule covered a wide variety of topics, including the woman's characteristics, marriage, fertility, contraception, children's immunizations and healthcare, nutrition, reproductive health, sexual behaviour, HIV/AIDS, women's empowerment, and domestic violence. The Man's Schedule covered the man's characteristics, marriage, his number of children, contraception, fertility preferences, nutrition, sexual behaviour, health issues, attitudes towards gender roles, and HIV/AIDS. The Biomarker Schedule covered measurements of height, weight, and haemoglobin levels for children; measurements of height, weight, waist and hip circumference, and haemoglobin levels for women age 15-49 years and men age 15-54 years; and blood pressure and random blood glucose levels for women and men age 15 years and over. In addition, women and men were requested to provide a few additional drops of blood from a finger prick for laboratory testing for HbA1c, malaria parasites, and Vitamin D3.

Readers should be cautious while interpreting and comparing the trends as some States/UTs may have smaller sample size. Moreover, at the time of survey, Ayushman Bharat AB-PMJAY and Pradhan Mantri Surakshit Matritva Abhiyan (PMSMA) were not fully rolled out and hence, their coverage may not have been factored in the results of indicator 12 (percentage of households with any usual member covered under a health insurance/financing scheme) and indicator 33 (percentage of mothers who received 4 or more antenatal care check-ups).

This fact sheet provides information on key indicators and trends for South Garo Hills. NFHS-5 fieldwork for Meghalaya was conducted from 8 July, 2019 to 15 November, 2019 by IQVIA Consulting and Information Services India Pvt. Ltd. In South Garo Hills, information was gathered from 921 households, 1,088 women, and 143 men.

South Garo Hills, Meghalaya - Key Indicators

Indicators	NFHS-5 (2019-20)	NFHS-4 (2015-16)
Population and Household Profile		
	Total	Total
1. Female population age 6 years and above who ever attended school (%)	89.2	86.2
2. Population below age 15 years (%)	31.4	31.4
3. Sex ratio of the total population (females per 1,000 males)	984	1,021
4. Sex ratio at birth for children born in the last five years (females per 1,000 males)	1,011	1,155
5. Children under age 5 years whose birth was registered with the civil authority (%)	92.0	94.5
6. Deaths in the last 3 years registered with the civil authority (%)	(53.8)	na
7. Population living in households with electricity (%)	98.6	99.0
8. Population living in households with an improved drinking-water source ¹ (%)	60.4	72.7
9. Population living in households that use an improved sanitation facility ² (%)	88.4	86.5
10. Households using clean fuel for cooking ³ (%)	22.3	9.8
11. Households using iodized salt (%)	83.6	98.9
12. Households with any usual member covered under a health insurance/financing scheme (%)	79.5	56.2
13. Children age 5 years who attended pre-primary school during the school year 2019-20 (%)	20.1	na
Characteristics of Women (age 15-49 years)		
14. Women who are literate ⁴ (%)	88.6	na
15. Women with 10 or more years of schooling (%)	38.4	51.5
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	10.2	10.2
17. Births in the 5 years preceding the survey that are third or higher order (%)	1.3	0.9
18. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	1.1	5.1
19. Women age 15-24 years who use hygienic methods of protection during their menstrual period ⁵ (%)	66.6	74.2
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ⁶ (%)	37.6	21.8
21. Any modern method ⁶ (%)	33.3	21.6
22. Female sterilization (%)	3.8	8.0
23. Male sterilization (%)	0.0	0.0
24. IUD/PPIUD (%)	9.0	3.0
25. Pill (%)	15.8	9.7
26. Condom (%)	2.8	0.5
27. Injectables (%)	1.6	0.4
Unmet Need for Family Planning (currently married women age 15-49 years)		
28. Total unmet need ⁷ (%)	25.3	16.2
29. Unmet need for spacing ⁷ (%)	14.4	11.7
Quality of Family Planning Services		
30. Health worker ever talked to female non-users about family planning (%)	20.3	42.4
31. Current users ever told about side effects of current method ⁸ (%)	63.7	(43.7)

Note: Major indicators are highlighted in grey.

LHV = Lady health visitor, ANM = Auxiliary nurse midwife

na = Not available

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rainwater, tanker truck, cart with small tank, bottled water, community RO plant.

²Flush to piped sewer system, flush to septic tank, flush to pit latrine, flush to don't know where, ventilated improved pit (VIP)/biogas latrine, pit latrine with slab, twin pit/composting toilet, which is not shared with any other household. This indicator does not denote access to toilet facility completely.

³Electricity, LPG/natural gas, biogas.

⁴Refers to women who completed standard 9 or higher and women who can read a whole sentence or part of a sentence.

⁵Locally prepared napkins, sanitary napkins, tampons, and menstrual cups are considered to be hygienic methods of protection.

⁶Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

⁷Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.
- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

Women who are classified as infecund have no unmet need because they are not at risk of becoming pregnant. Unmet need for family planning is the sum of unmet need for spacing plus unmet need for limiting.

⁸Based on current users of female sterilization, IUD/PPIUD, injectables, and pills who started using that method in the past 5 years.

South Garo Hills, Meghalaya - Key Indicators

Indicators	NFHS-5 (2019-20)	NFHS-4 (2015-16)
Maternal and Child Health	Total	Total
Maternity Care (for last birth in the 5 years before the survey)		
32. Mothers who had an antenatal check-up in the first trimester (%)	68.7	56.6
33. Mothers who had at least 4 antenatal care visits (%)	33.3	51.7
34. Mothers whose last birth was protected against neonatal tetanus ⁹ (%)	86.0	93.6
35. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	51.8	48.8
36. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	14.6	21.7
37. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	98.9	100.0
38. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery (%)	65.9	85.1
39. Average out-of-pocket expenditure per delivery in a public health facility (Rs.)	2,370	4,290
40. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	(0.0)	(0.0)
41. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery (%)	61.7	na
Delivery Care (for births in the 5 years before the survey)		
42. Institutional births (%)	88.3	88.4
43. Institutional births in public facility (%)	88.3	82.3
44. Home births that were conducted by skilled health personnel ¹⁰ (%)	3.9	0.5
45. Births attended by skilled health personnel ¹⁰ (%)	91.2	89.0
46. Births delivered by caesarean section (%)	4.7	8.1
47. Births in a private health facility that were delivered by caesarean section (%)	*	*
48. Births in a public health facility that were delivered by caesarean section (%)	5.3	8.0
Child Vaccinations and Vitamin A Supplementation		
49. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ¹¹ (%)	(86.8)	(83.0)
50. Children age 12-23 months fully vaccinated based on information from vaccination card only ¹² (%)	(89.1)	(88.4)
51. Children age 12-23 months who have received BCG (%)	(100.0)	(91.0)
52. Children age 12-23 months who have received 3 doses of polio vaccine ¹³ (%)	(94.7)	(93.9)
53. Children age 12-23 months who have received 3 doses of penta or DPT vaccine (%)	(86.8)	(93.9)
54. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	(89.6)	(85.1)
55. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	(15.0)	na
56. Children age 12-23 months who have received 3 doses of rotavirus vaccine ¹⁴ (%)	(2.4)	na
57. Children age 12-23 months who have received 3 doses of penta or hepatitis B vaccine (%)	(86.8)	(93.9)
58. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	57.2	78.4
59. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	(90.2)	(100.0)
60. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	(0.0)	(0.0)
Treatment of Childhood Diseases (children under age 5 years)		
61. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	5.4	28.6
62. Children with diarrhoea in the 2 weeks preceding the survey who received oral rehydration salts (ORS) (%)	*	97.7
63. Children with diarrhoea in the 2 weeks preceding the survey who received zinc (%)	*	76.0
64. Children with diarrhoea in the 2 weeks preceding the survey taken to a health facility or health provider (%)	*	64.7
65. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	3.1	14.1
66. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	(72.2)	75.0

⁹Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

¹⁰Doctor/nurse/LHV/ANM/midwife/other health personnel.

¹¹Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

¹²Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

¹³Not including polio vaccination given at birth.

¹⁴Since rotavirus is not being provided across all states and districts, the levels should not be compared.

South Garo Hills, Meghalaya - Key Indicators

Indicators	NFHS-5 (2019-20)	NFHS-4 (2015-16)
Child Feeding Practices and Nutritional Status of Children		
67. Children under age 3 years breastfed within one hour of birth ¹⁵ (%)	59.8	38.3
68. Children under age 6 months exclusively breastfed ¹⁶ (%)	*	(10.4)
69. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁶ (%)	*	*
70. Breastfeeding children age 6-23 months receiving an adequate diet ^{16, 17} (%)	23.0	36.5
71. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{16, 17} (%)	*	*
72. Total children age 6-23 months receiving an adequate diet ^{16, 17} (%)	23.0	37.4
73. Children under 5 years who are stunted (height-for-age) ¹⁸ (%)	29.8	16.8
74. Children under 5 years who are wasted (weight-for-height) ¹⁸ (%)	18.3	36.0
75. Children under 5 years who are severely wasted (weight-for-height) ¹⁹ (%)	8.8	23.7
76. Children under 5 years who are underweight (weight-for-age) ¹⁸ (%)	20.8	32.6
77. Children under 5 years who are overweight (weight-for-height) ²⁰ (%)	2.7	1.7
Nutritional Status of Women (age 15-49 years)		
78. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ²¹ (%)	6.6	3.4
79. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ²¹ (%)	6.9	14.2
80. Women who have high risk waist-to-hip ratio (≥0.85) (%)	67.7	na
Anaemia among Children and Women		
81. Children age 6-59 months who are anaemic (<11.0 g/dl) ²² (%)	33.6	74.9
82. Non-pregnant women age 15-49 years who are anaemic (<12.0 g/dl) ²² (%)	57.7	49.5
83. Pregnant women age 15-49 years who are anaemic (<11.0 g/dl) ²² (%)	33.7	27.9
84. All women age 15-49 years who are anaemic ²² (%)	56.2	46.6
85. All women age 15-19 years who are anaemic ²² (%)	65.0	35.2
Blood Sugar Level among Adults (age 15 years and above)		
Women		
86. Blood sugar level - high (141-160 mg/dl) ²³ (%)	10.0	na
87. Blood sugar level - very high (>160 mg/dl) ²³ (%)	2.1	na
88. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ²³ (%)	12.8	na
Men		
89. Blood sugar level - high (141-160 mg/dl) ²³ (%)	18.3	na
90. Blood sugar level - very high (>160 mg/dl) ²³ (%)	5.1	na
91. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ²³ (%)	23.7	na
Hypertension among Adults (age 15 years and above)		
Women		
92. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	10.1	na
93. Moderately or severely elevated blood pressure (Systolic ≥160mm of Hg and/or Diastolic ≥100mm of Hg) (%)	1.1	na
94. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	15.8	na
Men		
95. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	16.6	na
96. Moderately or severely elevated blood pressure (Systolic ≥160mm of Hg and/or Diastolic ≥100mm of Hg) (%)	3.3	na
97. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	22.8	na
Screening for Cancer among Women (age 30-49 years)		
98. Ever undergone a screening test for cervical cancer (%)	0.2	na
99. Ever undergone a breast examination for breast cancer (%)	0.0	na
100. Ever undergone an oral cavity examination for oral cancer (%)	0.0	na
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
101. Women age 15 years and above who use any kind of tobacco (%)	9.7	na
102. Men age 15 years and above who use any kind of tobacco (%)	50.7	na
103. Women age 15 years and above who consume alcohol (%)	1.4	na
104. Men age 15 years and above who consume alcohol (%)	33.1	na

¹⁵Based on the last child born in the 3 years before the survey.

¹⁶Based on the youngest child living with the mother.

¹⁷Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹⁸Below -2 standard deviations, based on the WHO standard.

¹⁹Below -3 standard deviations, based on the WHO standard.

²⁰Above +2 standard deviations, based on the WHO standard.

²¹Excludes pregnant women and women with a birth in the preceding 2 months.

²²Haemoglobin in grams per decilitre (g/dl). Among children, prevalence is adjusted for altitude. Among women, prevalence is adjusted for altitude and for smoking status, if known. As NFHS uses the capillary blood for estimation of anaemia, the results of NFHS-5 need not be compared with other surveys using venous blood.

²³Random blood sugar measurement.

NOTES



सत्यमेव जयते
Government of India

Ministry of Health and Family Welfare

NATIONAL FAMILY HEALTH SURVEY - 5

2019-20

DISTRICT FACT SHEET

SOUTH WEST GARO HILLS

MEGHALAYA



(स्थापना / Established in 1956)

बेहतर भविष्य के लिए क्षमता निर्माण
Capacity Building for a Better Future

International Institute for Population Sciences
(Deemed University)

Introduction

The National Family Health Survey 2019-20 (NFHS-5), the fifth in the NFHS series, provides information on population, health, and nutrition for India and each state/union territory (UT). Like NFHS-4, NFHS-5 also provides district-level estimates for many important indicators.

The contents of NFHS-5 are similar to NFHS-4 to allow comparisons over time. However, NFHS-5 includes some new topics, such as preschool education, disability, access to a toilet facility, death registration, bathing practices during menstruation, and methods and reasons for abortion. The scope of clinical, anthropometric, and biochemical testing (CAB) has also been expanded to include measurement of waist and hip circumferences, and the age range for the measurement of blood pressure and blood glucose has been expanded. However, HIV testing has been dropped. The NFHS-5 sample has been designed to provide national, state/union territory (UT), and district level estimates of various indicators covered in the survey. However, estimates of indicators of sexual behaviour; husband's background and woman's work; HIV/AIDS knowledge, attitudes and behaviour; and domestic violence are available only at the state/union territory (UT) and national level.

As in the earlier rounds, the Ministry of Health and Family Welfare, Government of India, designated the International Institute for Population Sciences, Mumbai, as the nodal agency to conduct NFHS-5. The main objective of each successive round of the NFHS has been to provide high-quality data on health and family welfare and emerging issues in this area. NFHS-5 data will be useful in setting benchmarks and examining the progress the health sector has made over time. Besides providing evidence for the effectiveness of ongoing programmes, the data from NFHS-5 help in identifying the need for new programmes with an area specific focus and identifying groups that are most in need of essential services.

Four Survey Schedules - Household, Woman's, Man's, and Biomarker - were canvassed in local languages using Computer Assisted Personal Interviewing (CAPI). In the Household Schedule, information was collected on all usual members of the household and visitors who stayed in the household the previous night, as well as socio-economic characteristics of the household; water, sanitation, and hygiene; health insurance coverage; disabilities; land ownership; number of deaths in the household in the three years preceding the survey; and the ownership and use of mosquito nets. The Woman's Schedule covered a wide variety of topics, including the woman's characteristics, marriage, fertility, contraception, children's immunizations and healthcare, nutrition, reproductive health, sexual behaviour, HIV/AIDS, women's empowerment, and domestic violence. The Man's Schedule covered the man's characteristics, marriage, his number of children, contraception, fertility preferences, nutrition, sexual behaviour, health issues, attitudes towards gender roles, and HIV/AIDS. The Biomarker Schedule covered measurements of height, weight, and haemoglobin levels for children; measurements of height, weight, waist and hip circumference, and haemoglobin levels for women age 15-49 years and men age 15-54 years; and blood pressure and random blood glucose levels for women and men age 15 years and over. In addition, women and men were requested to provide a few additional drops of blood from a finger prick for laboratory testing for HbA1c, malaria parasites, and Vitamin D3.

Readers should be cautious while interpreting and comparing the trends as some States/UTs may have smaller sample size. Moreover, at the time of survey, Ayushman Bharat AB-PMJAY and Pradhan Mantri Surakshit Matritva Abhiyan (PMSMA) were not fully rolled out and hence, their coverage may not have been factored in the results of indicator 12 (percentage of households with any usual member covered under a health insurance/financing scheme) and indicator 33 (percentage of mothers who received 4 or more antenatal care check-ups).

This fact sheet provides information on key indicators for South West Garo Hills. NFHS-5 fieldwork for Meghalaya was conducted from 8 July, 2019 to 15 November, 2019 by IQVIA Consulting and Information Services India Pvt. Ltd. In South West Garo Hills, information was gathered from 921 households, 1,161 women, and 140 men.

South West Garo Hills, Meghalaya - Key Indicators

Indicators	NFHS-5 (2019-20)
Population and Household Profile	
	Total
1. Female population age 6 years and above who ever attended school (%)	81.7
2. Population below age 15 years (%)	30.3
3. Sex ratio of the total population (females per 1,000 males)	995
4. Sex ratio at birth for children born in the last five years (females per 1,000 males)	1,036
5. Children under age 5 years whose birth was registered with the civil authority (%)	87.7
6. Deaths in the last 3 years registered with the civil authority (%)	71.4
7. Population living in households with electricity (%)	95.5
8. Population living in households with an improved drinking-water source ¹ (%)	58.6
9. Population living in households that use an improved sanitation facility ² (%)	71.4
10. Households using clean fuel for cooking ³ (%)	33.9
11. Households using iodized salt (%)	74.5
12. Households with any usual member covered under a health insurance/financing scheme (%)	75.1
13. Children age 5 years who attended pre-primary school during the school year 2019-20 (%)	7.8
Characteristics of Women (age 15-49 years)	
14. Women who are literate ⁴ (%)	82.6
15. Women with 10 or more years of schooling (%)	33.0
Marriage and Fertility	
16. Women age 20-24 years married before age 18 years (%)	15.3
17. Births in the 5 years preceding the survey that are third or higher order (%)	0.3
18. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	6.3
19. Women age 15-24 years who use hygienic methods of protection during their menstrual period ⁵ (%)	66.8
Current Use of Family Planning Methods (currently married women age 15-49 years)	
20. Any method ⁶ (%)	36.7
21. Any modern method ⁶ (%)	32.3
22. Female sterilization (%)	3.5
23. Male sterilization (%)	0.0
24. IUD/PPIUD (%)	5.5
25. Pill (%)	14.9
26. Condom (%)	6.2
27. Injectables (%)	1.0
Unmet Need for Family Planning (currently married women age 15-49 years)	
28. Total unmet need ⁷ (%)	30.4
29. Unmet need for spacing ⁷ (%)	16.7
Quality of Family Planning Services	
30. Health worker ever talked to female non-users about family planning (%)	24.2
31. Current users ever told about side effects of current method ⁸ (%)	82.9

Note: Indicator estimates for NFHS-4 are not shown in this table since no comparable estimates are available from NFHS-4 in this district due to district boundary changes or a newly formed district. Major indicators are highlighted in grey.

LHV = Lady health visitor, ANM = Auxiliary nurse midwife

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rainwater, tanker truck, cart with small tank, bottled water, community RO plant.

²Flush to piped sewer system, flush to septic tank, flush to pit latrine, flush to don't know where, ventilated improved pit (VIP)/biogas latrine, pit latrine with slab, twin pit/composting toilet, which is not shared with any other household. This indicator does not denote access to toilet facility completely.

³Electricity, LPG/natural gas, biogas.

⁴Refers to women who completed standard 9 or higher and women who can read a whole sentence or part of a sentence.

⁵Locally prepared napkins, sanitary napkins, tampons, and menstrual cups are considered to be hygienic methods of protection.

⁶Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

⁷Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.
- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

Women who are classified as infecund have no unmet need because they are not at risk of becoming pregnant. Unmet need for family planning is the sum of unmet need for spacing plus unmet need for limiting.

⁸Based on current users of female sterilization, IUD/PPIUD, injectables, and pills who started using that method in the past 5 years.

South West Garo Hills, Meghalaya - Key Indicators

Indicators	NFHS-5 (2019-20)
Maternal and Child Health	
Maternity Care (for last birth in the 5 years before the survey)	
32. Mothers who had an antenatal check-up in the first trimester (%)	57.0
33. Mothers who had at least 4 antenatal care visits (%)	28.5
34. Mothers whose last birth was protected against neonatal tetanus ⁹ (%)	80.7
35. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	28.3
36. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	11.0
37. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	99.0
38. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery (%)	60.6
39. Average out-of-pocket expenditure per delivery in a public health facility (Rs.)	2,666
40. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	(3.7)
41. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery (%)	55.1
Delivery Care (for births in the 5 years before the survey)	
42. Institutional births (%)	83.7
43. Institutional births in public facility (%)	83.7
44. Home births that were conducted by skilled health personnel ¹⁰ (%)	4.9
45. Births attended by skilled health personnel ¹⁰ (%)	88.3
46. Births delivered by caesarean section (%)	2.4
47. Births in a private health facility that were delivered by caesarean section (%)	*
48. Births in a public health facility that were delivered by caesarean section (%)	2.9
Child Vaccinations and Vitamin A Supplementation	
49. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ¹¹ (%)	85.4
50. Children age 12-23 months fully vaccinated based on information from vaccination card only ¹² (%)	90.1
51. Children age 12-23 months who have received BCG (%)	98.6
52. Children age 12-23 months who have received 3 doses of polio vaccine ¹³ (%)	91.1
53. Children age 12-23 months who have received 3 doses of penta or DPT vaccine (%)	90.9
54. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	92.1
55. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	25.7
56. Children age 12-23 months who have received 3 doses of rotavirus vaccine ¹⁴ (%)	13.4
57. Children age 12-23 months who have received 3 doses of penta or hepatitis B vaccine (%)	89.5
58. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	53.0
59. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	100.0
60. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	0.0
Treatment of Childhood Diseases (children under age 5 years)	
61. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	7.9
62. Children with diarrhoea in the 2 weeks preceding the survey who received oral rehydration salts (ORS) (%)	(89.5)
63. Children with diarrhoea in the 2 weeks preceding the survey who received zinc (%)	(46.4)
64. Children with diarrhoea in the 2 weeks preceding the survey taken to a health facility or health provider (%)	(85.9)
65. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	7.5
66. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	70.7

⁹Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

¹⁰Doctor/nurse/LHV/ANM/midwife/other health personnel.

¹¹Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

¹²Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

¹³Not including polio vaccination given at birth.

¹⁴Since rotavirus is not being provided across all states and districts, the levels should not be compared.

South West Garo Hills, Meghalaya - Key Indicators

Indicators	NFHS-5 (2019-20)
Child Feeding Practices and Nutritional Status of Children	
Total	
67. Children under age 3 years breastfed within one hour of birth ¹⁵ (%)	54.5
68. Children under age 6 months exclusively breastfed ¹⁶ (%)	*
69. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁶ (%)	*
70. Breastfeeding children age 6-23 months receiving an adequate diet ^{16, 17} (%)	32.1
71. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{16, 17} (%)	*
72. Total children age 6-23 months receiving an adequate diet ^{16, 17} (%)	30.0
73. Children under 5 years who are stunted (height-for-age) ¹⁸ (%)	32.2
74. Children under 5 years who are wasted (weight-for-height) ¹⁸ (%)	19.9
75. Children under 5 years who are severely wasted (weight-for-height) ¹⁹ (%)	7.1
76. Children under 5 years who are underweight (weight-for-age) ¹⁸ (%)	26.4
77. Children under 5 years who are overweight (weight-for-height) ²⁰ (%)	3.3
Nutritional Status of Women (age 15-49 years)	
78. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ²¹ (%)	8.5
79. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ²¹ (%)	9.0
80. Women who have high risk waist-to-hip ratio (≥0.85) (%)	71.5
Anaemia among Children and Women	
81. Children age 6-59 months who are anaemic (<11.0 g/dl) ²² (%)	35.5
82. Non-pregnant women age 15-49 years who are anaemic (<12.0 g/dl) ²² (%)	63.6
83. Pregnant women age 15-49 years who are anaemic (<11.0 g/dl) ²² (%)	37.8
84. All women age 15-49 years who are anaemic ²² (%)	62.5
85. All women age 15-19 years who are anaemic ²² (%)	67.3
Blood Sugar Level among Adults (age 15 years and above)	
Women	
86. Blood sugar level - high (141-160 mg/dl) ²³ (%)	8.9
87. Blood sugar level - very high (>160 mg/dl) ²³ (%)	3.5
88. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ²³ (%)	12.9
Men	
89. Blood sugar level - high (141-160 mg/dl) ²³ (%)	12.4
90. Blood sugar level - very high (>160 mg/dl) ²³ (%)	4.3
91. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ²³ (%)	16.7
Hypertension among Adults (age 15 years and above)	
Women	
92. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	8.0
93. Moderately or severely elevated blood pressure (Systolic ≥160mm of Hg and/or Diastolic ≥100mm of Hg) (%)	2.1
94. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	13.6
Men	
95. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	14.5
96. Moderately or severely elevated blood pressure (Systolic ≥160mm of Hg and/or Diastolic ≥100mm of Hg) (%)	2.7
97. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	20.0
Screening for Cancer among Women (age 30-49 years)	
98. Ever undergone a screening test for cervical cancer (%)	0.0
99. Ever undergone a breast examination for breast cancer (%)	0.0
100. Ever undergone an oral cavity examination for oral cancer (%)	0.0
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)	
101. Women age 15 years and above who use any kind of tobacco (%)	11.1
102. Men age 15 years and above who use any kind of tobacco (%)	52.4
103. Women age 15 years and above who consume alcohol (%)	1.2
104. Men age 15 years and above who consume alcohol (%)	30.6

¹⁵Based on the last child born in the 3 years before the survey.

¹⁶Based on the youngest child living with the mother.

¹⁷Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹⁸Below -2 standard deviations, based on the WHO standard.

¹⁹Below -3 standard deviations, based on the WHO standard.

²⁰Above +2 standard deviations, based on the WHO standard.

²¹Excludes pregnant women and women with a birth in the preceding 2 months.

²²Haemoglobin in grams per decilitre (g/dl). Among children, prevalence is adjusted for altitude. Among women, prevalence is adjusted for altitude and for smoking status, if known. As NFHS uses the capillary blood for estimation of anaemia, the results of NFHS-5 need not be compared with other surveys using venous blood.

²³Random blood sugar measurement.

NOTES



सत्यमेव जयते
Government of India

Ministry of Health and Family Welfare

NATIONAL FAMILY HEALTH SURVEY - 5

2019-20

DISTRICT FACT SHEET

SOUTH WEST KHASI HILLS

MEGHALAYA



(स्थापना / Established in 1956)

बेहतर भविष्य के लिए क्षमता निर्माण
Capacity Building for a Better Future

International Institute for Population Sciences
(Deemed University)

Introduction

The National Family Health Survey 2019-20 (NFHS-5), the fifth in the NFHS series, provides information on population, health, and nutrition for India and each state/union territory (UT). Like NFHS-4, NFHS-5 also provides district-level estimates for many important indicators.

The contents of NFHS-5 are similar to NFHS-4 to allow comparisons over time. However, NFHS-5 includes some new topics, such as preschool education, disability, access to a toilet facility, death registration, bathing practices during menstruation, and methods and reasons for abortion. The scope of clinical, anthropometric, and biochemical testing (CAB) has also been expanded to include measurement of waist and hip circumferences, and the age range for the measurement of blood pressure and blood glucose has been expanded. However, HIV testing has been dropped. The NFHS-5 sample has been designed to provide national, state/union territory (UT), and district level estimates of various indicators covered in the survey. However, estimates of indicators of sexual behaviour; husband's background and woman's work; HIV/AIDS knowledge, attitudes and behaviour; and domestic violence are available only at the state/union territory (UT) and national level.

As in the earlier rounds, the Ministry of Health and Family Welfare, Government of India, designated the International Institute for Population Sciences, Mumbai, as the nodal agency to conduct NFHS-5. The main objective of each successive round of the NFHS has been to provide high-quality data on health and family welfare and emerging issues in this area. NFHS-5 data will be useful in setting benchmarks and examining the progress the health sector has made over time. Besides providing evidence for the effectiveness of ongoing programmes, the data from NFHS-5 help in identifying the need for new programmes with an area specific focus and identifying groups that are most in need of essential services.

Four Survey Schedules - Household, Woman's, Man's, and Biomarker - were canvassed in local languages using Computer Assisted Personal Interviewing (CAPI). In the Household Schedule, information was collected on all usual members of the household and visitors who stayed in the household the previous night, as well as socio-economic characteristics of the household; water, sanitation, and hygiene; health insurance coverage; disabilities; land ownership; number of deaths in the household in the three years preceding the survey; and the ownership and use of mosquito nets. The Woman's Schedule covered a wide variety of topics, including the woman's characteristics, marriage, fertility, contraception, children's immunizations and healthcare, nutrition, reproductive health, sexual behaviour, HIV/AIDS, women's empowerment, and domestic violence. The Man's Schedule covered the man's characteristics, marriage, his number of children, contraception, fertility preferences, nutrition, sexual behaviour, health issues, attitudes towards gender roles, and HIV/AIDS. The Biomarker Schedule covered measurements of height, weight, and haemoglobin levels for children; measurements of height, weight, waist and hip circumference, and haemoglobin levels for women age 15-49 years and men age 15-54 years; and blood pressure and random blood glucose levels for women and men age 15 years and over. In addition, women and men were requested to provide a few additional drops of blood from a finger prick for laboratory testing for HbA1c, malaria parasites, and Vitamin D3.

Readers should be cautious while interpreting and comparing the trends as some States/UTs may have smaller sample size. Moreover, at the time of survey, Ayushman Bharat AB-PMJAY and Pradhan Mantri Surakshit Matritva Abhiyan (PMSMA) were not fully rolled out and hence, their coverage may not have been factored in the results of indicator 12 (percentage of households with any usual member covered under a health insurance/financing scheme) and indicator 33 (percentage of mothers who received 4 or more antenatal care check-ups).

This fact sheet provides information on key indicators for South West Khasi Hills. NFHS-5 fieldwork for Meghalaya was conducted from 8 July, 2019 to 15 November, 2019 by IQVIA Consulting and Information Services India Pvt. Ltd. In South West Khasi Hills, information was gathered from 924 households, 1,203 women, and 196 men.

South West Khasi Hills, Meghalaya - Key Indicators

Indicators	NFHS-5 (2019-20)
Population and Household Profile	
	Total
1. Female population age 6 years and above who ever attended school (%)	84.6
2. Population below age 15 years (%)	46.7
3. Sex ratio of the total population (females per 1,000 males)	1,025
4. Sex ratio at birth for children born in the last five years (females per 1,000 males)	1,032
5. Children under age 5 years whose birth was registered with the civil authority (%)	80.3
6. Deaths in the last 3 years registered with the civil authority (%)	49.5
7. Population living in households with electricity (%)	87.9
8. Population living in households with an improved drinking-water source ¹ (%)	85.3
9. Population living in households that use an improved sanitation facility ² (%)	89.6
10. Households using clean fuel for cooking ³ (%)	11.6
11. Households using iodized salt (%)	99.7
12. Households with any usual member covered under a health insurance/financing scheme (%)	72.3
13. Children age 5 years who attended pre-primary school during the school year 2019-20 (%)	44.2
Characteristics of Women (age 15-49 years)	
14. Women who are literate ⁴ (%)	85.9
15. Women with 10 or more years of schooling (%)	23.4
Marriage and Fertility	
16. Women age 20-24 years married before age 18 years (%)	21.6
17. Births in the 5 years preceding the survey that are third or higher order (%)	6.5
18. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	13.2
19. Women age 15-24 years who use hygienic methods of protection during their menstrual period ⁵ (%)	57.3
Current Use of Family Planning Methods (currently married women age 15-49 years)	
20. Any method ⁶ (%)	26.5
21. Any modern method ⁶ (%)	15.7
22. Female sterilization (%)	6.2
23. Male sterilization (%)	0.0
24. IUD/PPIUD (%)	1.2
25. Pill (%)	5.6
26. Condom (%)	1.2
27. Injectables (%)	1.1
Unmet Need for Family Planning (currently married women age 15-49 years)	
28. Total unmet need ⁷ (%)	26.2
29. Unmet need for spacing ⁷ (%)	20.1
Quality of Family Planning Services	
30. Health worker ever talked to female non-users about family planning (%)	42.8
31. Current users ever told about side effects of current method ⁸ (%)	82.3

Note: Indicator estimates for NFHS-4 are not shown in this table since no comparable estimates are available from NFHS-4 in this district due to district boundary changes or a newly formed district. Major indicators are highlighted in grey.

LHV = Lady health visitor, ANM = Auxiliary nurse midwife
() Based on 25-49 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rainwater, tanker truck, cart with small tank, bottled water, community RO plant.

²Flush to piped sewer system, flush to septic tank, flush to pit latrine, flush to don't know where, ventilated improved pit (VIP)/biogas latrine, pit latrine with slab, twin pit/composting toilet, which is not shared with any other household. This indicator does not denote access to toilet facility completely.

³Electricity, LPG/natural gas, biogas.

⁴Refers to women who completed standard 9 or higher and women who can read a whole sentence or part of a sentence.

⁵Locally prepared napkins, sanitary napkins, tampons, and menstrual cups are considered to be hygienic methods of protection.

⁶Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

⁷Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.
- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

Women who are classified as infecund have no unmet need because they are not at risk of becoming pregnant. Unmet need for family planning is the sum of unmet need for spacing plus unmet need for limiting.

⁸Based on current users of female sterilization, IUD/PPIUD, injectables, and pills who started using that method in the past 5 years.

South West Khasi Hills, Meghalaya - Key Indicators

Indicators	NFHS-5 (2019-20)
Maternal and Child Health	
Maternity Care (for last birth in the 5 years before the survey)	
32. Mothers who had an antenatal check-up in the first trimester (%)	48.5
33. Mothers who had at least 4 antenatal care visits (%)	63.4
34. Mothers whose last birth was protected against neonatal tetanus ⁹ (%)	85.6
35. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	53.8
36. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	19.8
37. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	97.0
38. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery (%)	42.9
39. Average out-of-pocket expenditure per delivery in a public health facility (Rs.)	3,902
40. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	0.3
41. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery (%)	38.7
Delivery Care (for births in the 5 years before the survey)	
42. Institutional births (%)	41.7
43. Institutional births in public facility (%)	37.6
44. Home births that were conducted by skilled health personnel ¹⁰ (%)	7.6
45. Births attended by skilled health personnel ¹⁰ (%)	49.3
46. Births delivered by caesarean section (%)	4.2
47. Births in a private health facility that were delivered by caesarean section (%)	(32.7)
48. Births in a public health facility that were delivered by caesarean section (%)	7.7
Child Vaccinations and Vitamin A Supplementation	
49. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ¹¹ (%)	78.8
50. Children age 12-23 months fully vaccinated based on information from vaccination card only ¹² (%)	86.3
51. Children age 12-23 months who have received BCG (%)	95.8
52. Children age 12-23 months who have received 3 doses of polio vaccine ¹³ (%)	80.5
53. Children age 12-23 months who have received 3 doses of penta or DPT vaccine (%)	86.9
54. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	86.4
55. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	17.4
56. Children age 12-23 months who have received 3 doses of rotavirus vaccine ¹⁴ (%)	0.6
57. Children age 12-23 months who have received 3 doses of penta or hepatitis B vaccine (%)	82.4
58. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	64.0
59. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	98.1
60. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	0.0
Treatment of Childhood Diseases (children under age 5 years)	
61. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	7.5
62. Children with diarrhoea in the 2 weeks preceding the survey who received oral rehydration salts (ORS) (%)	73.1
63. Children with diarrhoea in the 2 weeks preceding the survey who received zinc (%)	31.1
64. Children with diarrhoea in the 2 weeks preceding the survey taken to a health facility or health provider (%)	56.4
65. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	6.0
66. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	67.7

⁹Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

¹⁰Doctor/nurse/LHV/ANM/midwife/other health personnel.

¹¹Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

¹²Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

¹³Not including polio vaccination given at birth.

¹⁴Since rotavirus is not being provided across all states and districts, the levels should not be compared.

South West Khasi Hills, Meghalaya - Key Indicators

Indicators	NFHS-5 (2019-20)
Child Feeding Practices and Nutritional Status of Children	
Total	
67. Children under age 3 years breastfed within one hour of birth ¹⁵ (%)	76.8
68. Children under age 6 months exclusively breastfed ¹⁶ (%)	54.4
69. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁶ (%)	(65.6)
70. Breastfeeding children age 6-23 months receiving an adequate diet ^{16, 17} (%)	30.6
71. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{16, 17} (%)	33.5
72. Total children age 6-23 months receiving an adequate diet ^{16, 17} (%)	31.2
73. Children under 5 years who are stunted (height-for-age) ¹⁸ (%)	51.4
74. Children under 5 years who are wasted (weight-for-height) ¹⁸ (%)	10.4
75. Children under 5 years who are severely wasted (weight-for-height) ¹⁹ (%)	4.1
76. Children under 5 years who are underweight (weight-for-age) ¹⁸ (%)	27.6
77. Children under 5 years who are overweight (weight-for-height) ²⁰ (%)	3.5
Nutritional Status of Women (age 15-49 years)	
78. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ²¹ (%)	17.2
79. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ²¹ (%)	8.9
80. Women who have high risk waist-to-hip ratio (≥0.85) (%)	53.5
Anaemia among Children and Women	
81. Children age 6-59 months who are anaemic (<11.0 g/dl) ²² (%)	61.3
82. Non-pregnant women age 15-49 years who are anaemic (<12.0 g/dl) ²² (%)	58.5
83. Pregnant women age 15-49 years who are anaemic (<11.0 g/dl) ²² (%)	63.2
84. All women age 15-49 years who are anaemic ²² (%)	58.9
85. All women age 15-19 years who are anaemic ²² (%)	53.8
Blood Sugar Level among Adults (age 15 years and above)	
Women	
86. Blood sugar level - high (141-160 mg/dl) ²³ (%)	3.2
87. Blood sugar level - very high (>160 mg/dl) ²³ (%)	2.2
88. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ²³ (%)	6.3
Men	
89. Blood sugar level - high (141-160 mg/dl) ²³ (%)	4.2
90. Blood sugar level - very high (>160 mg/dl) ²³ (%)	2.6
91. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ²³ (%)	6.9
Hypertension among Adults (age 15 years and above)	
Women	
92. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	10.9
93. Moderately or severely elevated blood pressure (Systolic ≥160mm of Hg and/or Diastolic ≥100mm of Hg) (%)	4.4
94. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	18.5
Men	
95. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	12.8
96. Moderately or severely elevated blood pressure (Systolic ≥160mm of Hg and/or Diastolic ≥100mm of Hg) (%)	5.2
97. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	19.0
Screening for Cancer among Women (age 30-49 years)	
98. Ever undergone a screening test for cervical cancer (%)	0.2
99. Ever undergone a breast examination for breast cancer (%)	0.0
100. Ever undergone an oral cavity examination for oral cancer (%)	0.2
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)	
101. Women age 15 years and above who use any kind of tobacco (%)	50.0
102. Men age 15 years and above who use any kind of tobacco (%)	63.7
103. Women age 15 years and above who consume alcohol (%)	1.1
104. Men age 15 years and above who consume alcohol (%)	31.7

¹⁵Based on the last child born in the 3 years before the survey.

¹⁶Based on the youngest child living with the mother.

¹⁷Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹⁸Below -2 standard deviations, based on the WHO standard.

¹⁹Below -3 standard deviations, based on the WHO standard.

²⁰Above +2 standard deviations, based on the WHO standard.

²¹Excludes pregnant women and women with a birth in the preceding 2 months.

²²Haemoglobin in grams per decilitre (g/dl). Among children, prevalence is adjusted for altitude. Among women, prevalence is adjusted for altitude and for smoking status, if known. As NFHS uses the capillary blood for estimation of anaemia, the results of NFHS-5 need not be compared with other surveys using venous blood.

²³Random blood sugar measurement.

NOTES



सत्यमेव जयते
Government of India

Ministry of Health and Family Welfare

NATIONAL FAMILY HEALTH SURVEY - 5

2019-20

DISTRICT FACT SHEET

WEST GARO HILLS MEGHALAYA



(स्थापना / Established in 1956)

बेहतर भविष्य के लिए क्षमता निर्माण
Capacity Building for a Better Future

International Institute for Population Sciences
(Deemed University)

Introduction

The National Family Health Survey 2019-20 (NFHS-5), the fifth in the NFHS series, provides information on population, health, and nutrition for India and each state/union territory (UT). Like NFHS-4, NFHS-5 also provides district-level estimates for many important indicators.

The contents of NFHS-5 are similar to NFHS-4 to allow comparisons over time. However, NFHS-5 includes some new topics, such as preschool education, disability, access to a toilet facility, death registration, bathing practices during menstruation, and methods and reasons for abortion. The scope of clinical, anthropometric, and biochemical testing (CAB) has also been expanded to include measurement of waist and hip circumferences, and the age range for the measurement of blood pressure and blood glucose has been expanded. However, HIV testing has been dropped. The NFHS-5 sample has been designed to provide national, state/union territory (UT), and district level estimates of various indicators covered in the survey. However, estimates of indicators of sexual behaviour; husband's background and woman's work; HIV/AIDS knowledge, attitudes and behaviour; and domestic violence are available only at the state/union territory (UT) and national level.

As in the earlier rounds, the Ministry of Health and Family Welfare, Government of India, designated the International Institute for Population Sciences, Mumbai, as the nodal agency to conduct NFHS-5. The main objective of each successive round of the NFHS has been to provide high-quality data on health and family welfare and emerging issues in this area. NFHS-5 data will be useful in setting benchmarks and examining the progress the health sector has made over time. Besides providing evidence for the effectiveness of ongoing programmes, the data from NFHS-5 help in identifying the need for new programmes with an area specific focus and identifying groups that are most in need of essential services.

Four Survey Schedules - Household, Woman's, Man's, and Biomarker - were canvassed in local languages using Computer Assisted Personal Interviewing (CAPI). In the Household Schedule, information was collected on all usual members of the household and visitors who stayed in the household the previous night, as well as socio-economic characteristics of the household; water, sanitation, and hygiene; health insurance coverage; disabilities; land ownership; number of deaths in the household in the three years preceding the survey; and the ownership and use of mosquito nets. The Woman's Schedule covered a wide variety of topics, including the woman's characteristics, marriage, fertility, contraception, children's immunizations and healthcare, nutrition, reproductive health, sexual behaviour, HIV/AIDS, women's empowerment, and domestic violence. The Man's Schedule covered the man's characteristics, marriage, his number of children, contraception, fertility preferences, nutrition, sexual behaviour, health issues, attitudes towards gender roles, and HIV/AIDS. The Biomarker Schedule covered measurements of height, weight, and haemoglobin levels for children; measurements of height, weight, waist and hip circumference, and haemoglobin levels for women age 15-49 years and men age 15-54 years; and blood pressure and random blood glucose levels for women and men age 15 years and over. In addition, women and men were requested to provide a few additional drops of blood from a finger prick for laboratory testing for HbA1c, malaria parasites, and Vitamin D3.

Readers should be cautious while interpreting and comparing the trends as some States/UTs may have smaller sample size. Moreover, at the time of survey, Ayushman Bharat AB-PMJAY and Pradhan Mantri Surakshit Matritva Abhiyan (PMSMA) were not fully rolled out and hence, their coverage may not have been factored in the results of indicator 12 (percentage of households with any usual member covered under a health insurance/financing scheme) and indicator 33 (percentage of mothers who received 4 or more antenatal care check-ups).

This fact sheet provides information on key indicators for West Garo Hills. NFHS-5 fieldwork for Meghalaya was conducted from 8 July, 2019 to 15 November, 2019 by IQVIA Consulting and Information Services India Pvt. Ltd. In West Garo Hills, information was gathered from 923 households, 1,090 women, and 158 men.

West Garo Hills, Meghalaya - Key Indicators

Indicators	NFHS-5 (2019-20)
Population and Household Profile	
	Total
1. Female population age 6 years and above who ever attended school (%)	88.7
2. Population below age 15 years (%)	30.8
3. Sex ratio of the total population (females per 1,000 males)	1,025
4. Sex ratio at birth for children born in the last five years (females per 1,000 males)	1,396
5. Children under age 5 years whose birth was registered with the civil authority (%)	92.9
6. Deaths in the last 3 years registered with the civil authority (%)	(65.8)
7. Population living in households with electricity (%)	94.7
8. Population living in households with an improved drinking-water source ¹ (%)	83.7
9. Population living in households that use an improved sanitation facility ² (%)	86.3
10. Households using clean fuel for cooking ³ (%)	47.6
11. Households using iodized salt (%)	88.5
12. Households with any usual member covered under a health insurance/financing scheme (%)	77.9
13. Children age 5 years who attended pre-primary school during the school year 2019-20 (%)	20.2
Characteristics of Women (age 15-49 years)	
14. Women who are literate ⁴ (%)	89.0
15. Women with 10 or more years of schooling (%)	45.1
Marriage and Fertility	
16. Women age 20-24 years married before age 18 years (%)	10.4
17. Births in the 5 years preceding the survey that are third or higher order (%)	0.3
18. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	1.0
19. Women age 15-24 years who use hygienic methods of protection during their menstrual period ⁵ (%)	79.6
Current Use of Family Planning Methods (currently married women age 15-49 years)	
20. Any method ⁶ (%)	44.1
21. Any modern method ⁶ (%)	39.9
22. Female sterilization (%)	6.7
23. Male sterilization (%)	0.0
24. IUD/PPIUD (%)	9.5
25. Pill (%)	17.4
26. Condom (%)	4.1
27. Injectables (%)	1.6
Unmet Need for Family Planning (currently married women age 15-49 years)	
28. Total unmet need ⁷ (%)	19.7
29. Unmet need for spacing ⁷ (%)	9.2
Quality of Family Planning Services	
30. Health worker ever talked to female non-users about family planning (%)	32.0
31. Current users ever told about side effects of current method ⁸ (%)	58.1

Note: Indicator estimates for NFHS-4 are not shown in this table since no comparable estimates are available from NFHS-4 in this district due to district boundary changes or a newly formed district. Major indicators are highlighted in grey.

LHV = Lady health visitor, ANM = Auxiliary nurse midwife

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rainwater, tanker truck, cart with small tank, bottled water, community RO plant.

²Flush to piped sewer system, flush to septic tank, flush to pit latrine, flush to don't know where, ventilated improved pit (VIP)/biogas latrine, pit latrine with slab, twin pit/composting toilet, which is not shared with any other household. This indicator does not denote access to toilet facility completely.

³Electricity, LPG/natural gas, biogas.

⁴Refers to women who completed standard 9 or higher and women who can read a whole sentence or part of a sentence.

⁵Locally prepared napkins, sanitary napkins, tampons, and menstrual cups are considered to be hygienic methods of protection.

⁶Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

⁷Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.
- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

Women who are classified as infecund have no unmet need because they are not at risk of becoming pregnant. Unmet need for family planning is the sum of unmet need for spacing plus unmet need for limiting.

⁸Based on current users of female sterilization, IUD/PPIUD, injectables, and pills who started using that method in the past 5 years.

West Garo Hills, Meghalaya - Key Indicators

Indicators	NFHS-5 (2019-20)
Maternal and Child Health	Total
Maternity Care (for last birth in the 5 years before the survey)	
32. Mothers who had an antenatal check-up in the first trimester (%)	65.1
33. Mothers who had at least 4 antenatal care visits (%)	48.0
34. Mothers whose last birth was protected against neonatal tetanus ⁹ (%)	91.3
35. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	48.2
36. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	14.9
37. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	97.6
38. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery (%)	70.9
39. Average out-of-pocket expenditure per delivery in a public health facility (Rs.)	3,235
40. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	(2.1)
41. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery (%)	67.2
Delivery Care (for births in the 5 years before the survey)	
42. Institutional births (%)	89.6
43. Institutional births in public facility (%)	85.1
44. Home births that were conducted by skilled health personnel ¹⁰ (%)	3.2
45. Births attended by skilled health personnel ¹⁰ (%)	87.5
46. Births delivered by caesarean section (%)	8.4
47. Births in a private health facility that were delivered by caesarean section (%)	*
48. Births in a public health facility that were delivered by caesarean section (%)	8.2
Child Vaccinations and Vitamin A Supplementation	
49. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ¹¹ (%)	(77.1)
50. Children age 12-23 months fully vaccinated based on information from vaccination card only ¹² (%)	(78.6)
51. Children age 12-23 months who have received BCG (%)	(100.0)
52. Children age 12-23 months who have received 3 doses of polio vaccine ¹³ (%)	(84.1)
53. Children age 12-23 months who have received 3 doses of penta or DPT vaccine (%)	(84.8)
54. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	(87.4)
55. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	(5.3)
56. Children age 12-23 months who have received 3 doses of rotavirus vaccine ¹⁴ (%)	(5.0)
57. Children age 12-23 months who have received 3 doses of penta or hepatitis B vaccine (%)	(78.5)
58. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	68.2
59. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	(100.0)
60. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	(0.0)
Treatment of Childhood Diseases (children under age 5 years)	
61. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	7.8
62. Children with diarrhoea in the 2 weeks preceding the survey who received oral rehydration salts (ORS) (%)	*
63. Children with diarrhoea in the 2 weeks preceding the survey who received zinc (%)	*
64. Children with diarrhoea in the 2 weeks preceding the survey taken to a health facility or health provider (%)	*
65. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	3.5
66. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	(78.2)

⁹Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

¹⁰Doctor/nurse/LHV/ANM/midwife/other health personnel.

¹¹Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

¹²Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

¹³Not including polio vaccination given at birth.

¹⁴Since rotavirus is not being provided across all states and districts, the levels should not be compared.

West Garo Hills, Meghalaya - Key Indicators

Indicators	NFHS-5 (2019-20)
Child Feeding Practices and Nutritional Status of Children	
Total	
67. Children under age 3 years breastfed within one hour of birth ¹⁵ (%)	47.2
68. Children under age 6 months exclusively breastfed ¹⁶ (%)	*
69. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁶ (%)	*
70. Breastfeeding children age 6-23 months receiving an adequate diet ^{16, 17} (%)	14.6
71. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{16, 17} (%)	*
72. Total children age 6-23 months receiving an adequate diet ^{16, 17} (%)	17.2
73. Children under 5 years who are stunted (height-for-age) ¹⁸ (%)	39.5
74. Children under 5 years who are wasted (weight-for-height) ¹⁸ (%)	14.9
75. Children under 5 years who are severely wasted (weight-for-height) ¹⁹ (%)	4.7
76. Children under 5 years who are underweight (weight-for-age) ¹⁸ (%)	26.3
77. Children under 5 years who are overweight (weight-for-height) ²⁰ (%)	5.2
Nutritional Status of Women (age 15-49 years)	
78. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ²¹ (%)	7.7
79. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ²¹ (%)	7.2
80. Women who have high risk waist-to-hip ratio (≥0.85) (%)	74.0
Anaemia among Children and Women	
81. Children age 6-59 months who are anaemic (<11.0 g/dl) ²² (%)	30.2
82. Non-pregnant women age 15-49 years who are anaemic (<12.0 g/dl) ²² (%)	57.2
83. Pregnant women age 15-49 years who are anaemic (<11.0 g/dl) ²² (%)	26.7
84. All women age 15-49 years who are anaemic ²² (%)	55.3
85. All women age 15-19 years who are anaemic ²² (%)	62.0
Blood Sugar Level among Adults (age 15 years and above)	
Women	
86. Blood sugar level - high (141-160 mg/dl) ²³ (%)	7.6
87. Blood sugar level - very high (>160 mg/dl) ²³ (%)	3.3
88. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ²³ (%)	12.1
Men	
89. Blood sugar level - high (141-160 mg/dl) ²³ (%)	15.9
90. Blood sugar level - very high (>160 mg/dl) ²³ (%)	7.9
91. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ²³ (%)	24.6
Hypertension among Adults (age 15 years and above)	
Women	
92. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	11.5
93. Moderately or severely elevated blood pressure (Systolic ≥160mm of Hg and/or Diastolic ≥100mm of Hg) (%)	2.7
94. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	17.4
Men	
95. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	17.2
96. Moderately or severely elevated blood pressure (Systolic ≥160mm of Hg and/or Diastolic ≥100mm of Hg) (%)	3.4
97. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	24.7
Screening for Cancer among Women (age 30-49 years)	
98. Ever undergone a screening test for cervical cancer (%)	0.0
99. Ever undergone a breast examination for breast cancer (%)	0.2
100. Ever undergone an oral cavity examination for oral cancer (%)	0.0
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)	
101. Women age 15 years and above who use any kind of tobacco (%)	17.6
102. Men age 15 years and above who use any kind of tobacco (%)	58.3
103. Women age 15 years and above who consume alcohol (%)	2.1
104. Men age 15 years and above who consume alcohol (%)	38.8

¹⁵Based on the last child born in the 3 years before the survey.

¹⁶Based on the youngest child living with the mother.

¹⁷Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹⁸Below -2 standard deviations, based on the WHO standard.

¹⁹Below -3 standard deviations, based on the WHO standard.

²⁰Above +2 standard deviations, based on the WHO standard.

²¹Excludes pregnant women and women with a birth in the preceding 2 months.

²²Haemoglobin in grams per decilitre (g/dl). Among children, prevalence is adjusted for altitude. Among women, prevalence is adjusted for altitude and for smoking status, if known. As NFHS uses the capillary blood for estimation of anaemia, the results of NFHS-5 need not be compared with other surveys using venous blood.

²³Random blood sugar measurement.

NOTES



सत्यमेव जयते
Government of India

Ministry of Health and Family Welfare

NATIONAL FAMILY HEALTH SURVEY - 5

2019-20

DISTRICT FACT SHEET

WEST JAIINTIA HILLS MEGHALAYA



(स्थापना / Established in 1956)

बेहतर भविष्य के लिए क्षमता निर्माण
Capacity Building for a Better Future

International Institute for Population Sciences
(Deemed University)

Introduction

The National Family Health Survey 2019-20 (NFHS-5), the fifth in the NFHS series, provides information on population, health, and nutrition for India and each state/union territory (UT). Like NFHS-4, NFHS-5 also provides district-level estimates for many important indicators.

The contents of NFHS-5 are similar to NFHS-4 to allow comparisons over time. However, NFHS-5 includes some new topics, such as preschool education, disability, access to a toilet facility, death registration, bathing practices during menstruation, and methods and reasons for abortion. The scope of clinical, anthropometric, and biochemical testing (CAB) has also been expanded to include measurement of waist and hip circumferences, and the age range for the measurement of blood pressure and blood glucose has been expanded. However, HIV testing has been dropped. The NFHS-5 sample has been designed to provide national, state/union territory (UT), and district level estimates of various indicators covered in the survey. However, estimates of indicators of sexual behaviour; husband's background and woman's work; HIV/AIDS knowledge, attitudes and behaviour; and domestic violence are available only at the state/union territory (UT) and national level.

As in the earlier rounds, the Ministry of Health and Family Welfare, Government of India, designated the International Institute for Population Sciences, Mumbai, as the nodal agency to conduct NFHS-5. The main objective of each successive round of the NFHS has been to provide high-quality data on health and family welfare and emerging issues in this area. NFHS-5 data will be useful in setting benchmarks and examining the progress the health sector has made over time. Besides providing evidence for the effectiveness of ongoing programmes, the data from NFHS-5 help in identifying the need for new programmes with an area specific focus and identifying groups that are most in need of essential services.

Four Survey Schedules - Household, Woman's, Man's, and Biomarker - were canvassed in local languages using Computer Assisted Personal Interviewing (CAPI). In the Household Schedule, information was collected on all usual members of the household and visitors who stayed in the household the previous night, as well as socio-economic characteristics of the household; water, sanitation, and hygiene; health insurance coverage; disabilities; land ownership; number of deaths in the household in the three years preceding the survey; and the ownership and use of mosquito nets. The Woman's Schedule covered a wide variety of topics, including the woman's characteristics, marriage, fertility, contraception, children's immunizations and healthcare, nutrition, reproductive health, sexual behaviour, HIV/AIDS, women's empowerment, and domestic violence. The Man's Schedule covered the man's characteristics, marriage, his number of children, contraception, fertility preferences, nutrition, sexual behaviour, health issues, attitudes towards gender roles, and HIV/AIDS. The Biomarker Schedule covered measurements of height, weight, and haemoglobin levels for children; measurements of height, weight, waist and hip circumference, and haemoglobin levels for women age 15-49 years and men age 15-54 years; and blood pressure and random blood glucose levels for women and men age 15 years and over. In addition, women and men were requested to provide a few additional drops of blood from a finger prick for laboratory testing for HbA1c, malaria parasites, and Vitamin D3.

Readers should be cautious while interpreting and comparing the trends as some States/UTs may have smaller sample size. Moreover, at the time of survey, Ayushman Bharat AB-PMJAY and Pradhan Mantri Surakshit Matritva Abhiyan (PMSMA) were not fully rolled out and hence, their coverage may not have been factored in the results of indicator 12 (percentage of households with any usual member covered under a health insurance/financing scheme) and indicator 33 (percentage of mothers who received 4 or more antenatal care check-ups).

This fact sheet provides information on key indicators for West Jaintia Hills. NFHS-5 fieldwork for Meghalaya was conducted from 8 July, 2019 to 15 November, 2019 by IQVIA Consulting and Information Services India Pvt. Ltd. In West Jaintia Hills, information was gathered from 924 households, 1,172 women, and 155 men.

West Jaintia Hills, Meghalaya - Key Indicators

Indicators	NFHS-5 (2019-20)
Population and Household Profile	
	Total
1. Female population age 6 years and above who ever attended school (%)	73.8
2. Population below age 15 years (%)	44.8
3. Sex ratio of the total population (females per 1,000 males)	1,143
4. Sex ratio at birth for children born in the last five years (females per 1,000 males)	1,036
5. Children under age 5 years whose birth was registered with the civil authority (%)	81.7
6. Deaths in the last 3 years registered with the civil authority (%)	55.7
7. Population living in households with electricity (%)	75.4
8. Population living in households with an improved drinking-water source ¹ (%)	70.6
9. Population living in households that use an improved sanitation facility ² (%)	75.1
10. Households using clean fuel for cooking ³ (%)	20.7
11. Households using iodized salt (%)	97.3
12. Households with any usual member covered under a health insurance/financing scheme (%)	65.6
13. Children age 5 years who attended pre-primary school during the school year 2019-20 (%)	33.3
Characteristics of Women (age 15-49 years)	
14. Women who are literate ⁴ (%)	80.0
15. Women with 10 or more years of schooling (%)	24.7
Marriage and Fertility	
16. Women age 20-24 years married before age 18 years (%)	19.9
17. Births in the 5 years preceding the survey that are third or higher order (%)	4.8
18. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	11.9
19. Women age 15-24 years who use hygienic methods of protection during their menstrual period ⁵ (%)	32.6
Current Use of Family Planning Methods (currently married women age 15-49 years)	
20. Any method ⁶ (%)	19.3
21. Any modern method ⁶ (%)	17.5
22. Female sterilization (%)	4.5
23. Male sterilization (%)	0.0
24. IUD/PPIUD (%)	2.5
25. Pill (%)	7.1
26. Condom (%)	1.0
27. Injectables (%)	2.4
Unmet Need for Family Planning (currently married women age 15-49 years)	
28. Total unmet need ⁷ (%)	29.3
29. Unmet need for spacing ⁷ (%)	21.7
Quality of Family Planning Services	
30. Health worker ever talked to female non-users about family planning (%)	31.4
31. Current users ever told about side effects of current method ⁸ (%)	70.4

Note: Indicator estimates for NFHS-4 are not shown in this table since no comparable estimates are available from NFHS-4 in this district due to district boundary changes or a newly formed district. Major indicators are highlighted in grey.

LHV = Lady health visitor, ANM = Auxiliary nurse midwife
() Based on 25-49 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rainwater, tanker truck, cart with small tank, bottled water, community RO plant.

²Flush to piped sewer system, flush to septic tank, flush to pit latrine, flush to don't know where, ventilated improved pit (VIP)/biogas latrine, pit latrine with slab, twin pit/composting toilet, which is not shared with any other household. This indicator does not denote access to toilet facility completely.

³Electricity, LPG/natural gas, biogas.

⁴Refers to women who completed standard 9 or higher and women who can read a whole sentence or part of a sentence.

⁵Locally prepared napkins, sanitary napkins, tampons, and menstrual cups are considered to be hygienic methods of protection.

⁶Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

⁷Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.
- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

Women who are classified as infecund have no unmet need because they are not at risk of becoming pregnant. Unmet need for family planning is the sum of unmet need for spacing plus unmet need for limiting.

⁸Based on current users of female sterilization, IUD/PPIUD, injectables, and pills who started using that method in the past 5 years.

West Jaintia Hills, Meghalaya - Key Indicators

Indicators	NFHS-5 (2019-20)
Maternal and Child Health	
Maternity Care (for last birth in the 5 years before the survey)	
32. Mothers who had an antenatal check-up in the first trimester (%)	44.0
33. Mothers who had at least 4 antenatal care visits (%)	43.1
34. Mothers whose last birth was protected against neonatal tetanus ⁹ (%)	79.2
35. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	32.7
36. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	19.8
37. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	88.1
38. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery (%)	28.6
39. Average out-of-pocket expenditure per delivery in a public health facility (Rs.)	2,917
40. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	0.4
41. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery (%)	30.8
Delivery Care (for births in the 5 years before the survey)	
42. Institutional births (%)	42.2
43. Institutional births in public facility (%)	31.9
44. Home births that were conducted by skilled health personnel ¹⁰ (%)	7.7
45. Births attended by skilled health personnel ¹⁰ (%)	50.7
46. Births delivered by caesarean section (%)	6.2
47. Births in a private health facility that were delivered by caesarean section (%)	34.4
48. Births in a public health facility that were delivered by caesarean section (%)	8.3
Child Vaccinations and Vitamin A Supplementation	
49. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ¹¹ (%)	59.7
50. Children age 12-23 months fully vaccinated based on information from vaccination card only ¹² (%)	84.1
51. Children age 12-23 months who have received BCG (%)	90.9
52. Children age 12-23 months who have received 3 doses of polio vaccine ¹³ (%)	65.2
53. Children age 12-23 months who have received 3 doses of penta or DPT vaccine (%)	77.0
54. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	72.3
55. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	22.4
56. Children age 12-23 months who have received 3 doses of rotavirus vaccine ¹⁴ (%)	4.8
57. Children age 12-23 months who have received 3 doses of penta or hepatitis B vaccine (%)	72.0
58. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	63.2
59. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	93.3
60. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	2.3
Treatment of Childhood Diseases (children under age 5 years)	
61. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	13.7
62. Children with diarrhoea in the 2 weeks preceding the survey who received oral rehydration salts (ORS) (%)	71.3
63. Children with diarrhoea in the 2 weeks preceding the survey who received zinc (%)	41.9
64. Children with diarrhoea in the 2 weeks preceding the survey taken to a health facility or health provider (%)	67.6
65. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	4.8
66. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	79.1

⁹Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

¹⁰Doctor/nurse/LHV/ANM/midwife/other health personnel.

¹¹Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

¹²Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

¹³Not including polio vaccination given at birth.

¹⁴Since rotavirus is not being provided across all states and districts, the levels should not be compared.

West Jaintia Hills, Meghalaya - Key Indicators

Indicators	NFHS-5 (2019-20)
Child Feeding Practices and Nutritional Status of Children	
Total	
67. Children under age 3 years breastfed within one hour of birth ¹⁵ (%)	88.5
68. Children under age 6 months exclusively breastfed ¹⁶ (%)	46.4
69. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁶ (%)	(77.7)
70. Breastfeeding children age 6-23 months receiving an adequate diet ^{16, 17} (%)	25.7
71. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{16, 17} (%)	20.8
72. Total children age 6-23 months receiving an adequate diet ^{16, 17} (%)	24.5
73. Children under 5 years who are stunted (height-for-age) ¹⁸ (%)	48.7
74. Children under 5 years who are wasted (weight-for-height) ¹⁸ (%)	9.7
75. Children under 5 years who are severely wasted (weight-for-height) ¹⁹ (%)	4.0
76. Children under 5 years who are underweight (weight-for-age) ¹⁸ (%)	28.3
77. Children under 5 years who are overweight (weight-for-height) ²⁰ (%)	3.0
Nutritional Status of Women (age 15-49 years)	
78. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ²¹ (%)	9.6
79. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ²¹ (%)	15.4
80. Women who have high risk waist-to-hip ratio (≥0.85) (%)	59.1
Anaemia among Children and Women	
81. Children age 6-59 months who are anaemic (<11.0 g/dl) ²² (%)	54.6
82. Non-pregnant women age 15-49 years who are anaemic (<12.0 g/dl) ²² (%)	55.3
83. Pregnant women age 15-49 years who are anaemic (<11.0 g/dl) ²² (%)	64.6
84. All women age 15-49 years who are anaemic ²² (%)	55.9
85. All women age 15-19 years who are anaemic ²² (%)	54.2
Blood Sugar Level among Adults (age 15 years and above)	
Women	
86. Blood sugar level - high (141-160 mg/dl) ²³ (%)	3.8
87. Blood sugar level - very high (>160 mg/dl) ²³ (%)	3.5
88. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ²³ (%)	8.8
Men	
89. Blood sugar level - high (141-160 mg/dl) ²³ (%)	4.4
90. Blood sugar level - very high (>160 mg/dl) ²³ (%)	2.8
91. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ²³ (%)	7.8
Hypertension among Adults (age 15 years and above)	
Women	
92. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	9.0
93. Moderately or severely elevated blood pressure (Systolic ≥160mm of Hg and/or Diastolic ≥100mm of Hg) (%)	3.3
94. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	18.0
Men	
95. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	8.6
96. Moderately or severely elevated blood pressure (Systolic ≥160mm of Hg and/or Diastolic ≥100mm of Hg) (%)	3.2
97. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	15.5
Screening for Cancer among Women (age 30-49 years)	
98. Ever undergone a screening test for cervical cancer (%)	0.5
99. Ever undergone a breast examination for breast cancer (%)	0.1
100. Ever undergone an oral cavity examination for oral cancer (%)	0.4
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)	
101. Women age 15 years and above who use any kind of tobacco (%)	44.8
102. Men age 15 years and above who use any kind of tobacco (%)	64.7
103. Women age 15 years and above who consume alcohol (%)	1.3
104. Men age 15 years and above who consume alcohol (%)	36.9

¹⁵Based on the last child born in the 3 years before the survey.

¹⁶Based on the youngest child living with the mother.

¹⁷Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹⁸Below -2 standard deviations, based on the WHO standard.

¹⁹Below -3 standard deviations, based on the WHO standard.

²⁰Above +2 standard deviations, based on the WHO standard.

²¹Excludes pregnant women and women with a birth in the preceding 2 months.

²²Haemoglobin in grams per decilitre (g/dl). Among children, prevalence is adjusted for altitude. Among women, prevalence is adjusted for altitude and for smoking status, if known. As NFHS uses the capillary blood for estimation of anaemia, the results of NFHS-5 need not be compared with other surveys using venous blood.

²³Random blood sugar measurement.

NOTES



सत्यमेव जयते
Government of India

Ministry of Health and Family Welfare

NATIONAL FAMILY HEALTH SURVEY - 5

2019-20

DISTRICT FACT SHEET

WEST KHASI HILLS MEGHALAYA



(स्थापना / Established in 1956)

बेहतर भविष्य के लिए क्षमता निर्माण
Capacity Building for a Better Future

International Institute for Population Sciences
(Deemed University)

Introduction

The National Family Health Survey 2019-20 (NFHS-5), the fifth in the NFHS series, provides information on population, health, and nutrition for India and each state/union territory (UT). Like NFHS-4, NFHS-5 also provides district-level estimates for many important indicators.

The contents of NFHS-5 are similar to NFHS-4 to allow comparisons over time. However, NFHS-5 includes some new topics, such as preschool education, disability, access to a toilet facility, death registration, bathing practices during menstruation, and methods and reasons for abortion. The scope of clinical, anthropometric, and biochemical testing (CAB) has also been expanded to include measurement of waist and hip circumferences, and the age range for the measurement of blood pressure and blood glucose has been expanded. However, HIV testing has been dropped. The NFHS-5 sample has been designed to provide national, state/union territory (UT), and district level estimates of various indicators covered in the survey. However, estimates of indicators of sexual behaviour; husband's background and woman's work; HIV/AIDS knowledge, attitudes and behaviour; and domestic violence are available only at the state/union territory (UT) and national level.

As in the earlier rounds, the Ministry of Health and Family Welfare, Government of India, designated the International Institute for Population Sciences, Mumbai, as the nodal agency to conduct NFHS-5. The main objective of each successive round of the NFHS has been to provide high-quality data on health and family welfare and emerging issues in this area. NFHS-5 data will be useful in setting benchmarks and examining the progress the health sector has made over time. Besides providing evidence for the effectiveness of ongoing programmes, the data from NFHS-5 help in identifying the need for new programmes with an area specific focus and identifying groups that are most in need of essential services.

Four Survey Schedules - Household, Woman's, Man's, and Biomarker - were canvassed in local languages using Computer Assisted Personal Interviewing (CAPI). In the Household Schedule, information was collected on all usual members of the household and visitors who stayed in the household the previous night, as well as socio-economic characteristics of the household; water, sanitation, and hygiene; health insurance coverage; disabilities; land ownership; number of deaths in the household in the three years preceding the survey; and the ownership and use of mosquito nets. The Woman's Schedule covered a wide variety of topics, including the woman's characteristics, marriage, fertility, contraception, children's immunizations and healthcare, nutrition, reproductive health, sexual behaviour, HIV/AIDS, women's empowerment, and domestic violence. The Man's Schedule covered the man's characteristics, marriage, his number of children, contraception, fertility preferences, nutrition, sexual behaviour, health issues, attitudes towards gender roles, and HIV/AIDS. The Biomarker Schedule covered measurements of height, weight, and haemoglobin levels for children; measurements of height, weight, waist and hip circumference, and haemoglobin levels for women age 15-49 years and men age 15-54 years; and blood pressure and random blood glucose levels for women and men age 15 years and over. In addition, women and men were requested to provide a few additional drops of blood from a finger prick for laboratory testing for HbA1c, malaria parasites, and Vitamin D3.

Readers should be cautious while interpreting and comparing the trends as some States/UTs may have smaller sample size. Moreover, at the time of survey, Ayushman Bharat AB-PMJAY and Pradhan Mantri Surakshit Matritva Abhiyan (PMSMA) were not fully rolled out and hence, their coverage may not have been factored in the results of indicator 12 (percentage of households with any usual member covered under a health insurance/financing scheme) and indicator 33 (percentage of mothers who received 4 or more antenatal care check-ups).

This fact sheet provides information on key indicators for West Khasi Hills. NFHS-5 fieldwork for Meghalaya was conducted from 8 July, 2019 to 15 November, 2019 by IQVIA Consulting and Information Services India Pvt. Ltd. In West Khasi Hills, information was gathered from 924 households, 1,197 women, and 178 men.

West Khasi Hills, Meghalaya - Key Indicators

Indicators	NFHS-5 (2019-20)
Population and Household Profile	
	Total
1. Female population age 6 years and above who ever attended school (%)	85.1
2. Population below age 15 years (%)	50.6
3. Sex ratio of the total population (females per 1,000 males)	1,038
4. Sex ratio at birth for children born in the last five years (females per 1,000 males)	984
5. Children under age 5 years whose birth was registered with the civil authority (%)	79.2
6. Deaths in the last 3 years registered with the civil authority (%)	49.0
7. Population living in households with electricity (%)	85.0
8. Population living in households with an improved drinking-water source ¹ (%)	80.5
9. Population living in households that use an improved sanitation facility ² (%)	87.8
10. Households using clean fuel for cooking ³ (%)	8.6
11. Households using iodized salt (%)	97.3
12. Households with any usual member covered under a health insurance/financing scheme (%)	60.1
13. Children age 5 years who attended pre-primary school during the school year 2019-20 (%)	34.8
Characteristics of Women (age 15-49 years)	
14. Women who are literate ⁴ (%)	87.4
15. Women with 10 or more years of schooling (%)	16.4
Marriage and Fertility	
16. Women age 20-24 years married before age 18 years (%)	30.3
17. Births in the 5 years preceding the survey that are third or higher order (%)	8.0
18. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	12.9
19. Women age 15-24 years who use hygienic methods of protection during their menstrual period ⁵ (%)	48.9
Current Use of Family Planning Methods (currently married women age 15-49 years)	
20. Any method ⁶ (%)	26.3
21. Any modern method ⁶ (%)	15.3
22. Female sterilization (%)	9.3
23. Male sterilization (%)	0.0
24. IUD/PPIUD (%)	1.6
25. Pill (%)	2.8
26. Condom (%)	0.5
27. Injectables (%)	1.1
Unmet Need for Family Planning (currently married women age 15-49 years)	
28. Total unmet need ⁷ (%)	25.8
29. Unmet need for spacing ⁷ (%)	21.1
Quality of Family Planning Services	
30. Health worker ever talked to female non-users about family planning (%)	43.4
31. Current users ever told about side effects of current method ⁸ (%)	72.8

Note: Indicator estimates for NFHS-4 are not shown in this table since no comparable estimates are available from NFHS-4 in this district due to district boundary changes or a newly formed district. Major indicators are highlighted in grey.

LHV = Lady health visitor, ANM = Auxiliary nurse midwife
() Based on 25-49 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rainwater, tanker truck, cart with small tank, bottled water, community RO plant.

²Flush to piped sewer system, flush to septic tank, flush to pit latrine, flush to don't know where, ventilated improved pit (VIP)/biogas latrine, pit latrine with slab, twin pit/composting toilet, which is not shared with any other household. This indicator does not denote access to toilet facility completely.

³Electricity, LPG/natural gas, biogas.

⁴Refers to women who completed standard 9 or higher and women who can read a whole sentence or part of a sentence.

⁵Locally prepared napkins, sanitary napkins, tampons, and menstrual cups are considered to be hygienic methods of protection.

⁶Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

⁷Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.
- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

Women who are classified as infecund have no unmet need because they are not at risk of becoming pregnant. Unmet need for family planning is the sum of unmet need for spacing plus unmet need for limiting.

⁸Based on current users of female sterilization, IUD/PPIUD, injectables, and pills who started using that method in the past 5 years.

West Khasi Hills, Meghalaya - Key Indicators

Indicators	NFHS-5 (2019-20)
Maternal and Child Health	
Maternity Care (for last birth in the 5 years before the survey)	
32. Mothers who had an antenatal check-up in the first trimester (%)	45.3
33. Mothers who had at least 4 antenatal care visits (%)	64.0
34. Mothers whose last birth was protected against neonatal tetanus ⁹ (%)	81.6
35. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	51.6
36. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	21.0
37. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	97.3
38. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery (%)	42.0
39. Average out-of-pocket expenditure per delivery in a public health facility (Rs.)	2,947
40. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	0.4
41. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery (%)	41.4
Delivery Care (for births in the 5 years before the survey)	
42. Institutional births (%)	41.7
43. Institutional births in public facility (%)	36.1
44. Home births that were conducted by skilled health personnel ¹⁰ (%)	7.2
45. Births attended by skilled health personnel ¹⁰ (%)	49.3
46. Births delivered by caesarean section (%)	4.7
47. Births in a private health facility that were delivered by caesarean section (%)	30.6
48. Births in a public health facility that were delivered by caesarean section (%)	8.3
Child Vaccinations and Vitamin A Supplementation	
49. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ¹¹ (%)	62.5
50. Children age 12-23 months fully vaccinated based on information from vaccination card only ¹² (%)	77.4
51. Children age 12-23 months who have received BCG (%)	86.7
52. Children age 12-23 months who have received 3 doses of polio vaccine ¹³ (%)	67.6
53. Children age 12-23 months who have received 3 doses of penta or DPT vaccine (%)	68.2
54. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	71.5
55. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	12.3
56. Children age 12-23 months who have received 3 doses of rotavirus vaccine ¹⁴ (%)	0.7
57. Children age 12-23 months who have received 3 doses of penta or hepatitis B vaccine (%)	66.4
58. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	63.4
59. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	95.9
60. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	0.9
Treatment of Childhood Diseases (children under age 5 years)	
61. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	10.1
62. Children with diarrhoea in the 2 weeks preceding the survey who received oral rehydration salts (ORS) (%)	75.1
63. Children with diarrhoea in the 2 weeks preceding the survey who received zinc (%)	48.8
64. Children with diarrhoea in the 2 weeks preceding the survey taken to a health facility or health provider (%)	77.4
65. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	6.5
66. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	75.3

⁹Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

¹⁰Doctor/nurse/LHV/ANM/midwife/other health personnel.

¹¹Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

¹²Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

¹³Not including polio vaccination given at birth.

¹⁴Since rotavirus is not being provided across all states and districts, the levels should not be compared.

West Khasi Hills, Meghalaya - Key Indicators

Indicators	NFHS-5 (2019-20)
Child Feeding Practices and Nutritional Status of Children	
Total	
67. Children under age 3 years breastfed within one hour of birth ¹⁵ (%)	81.2
68. Children under age 6 months exclusively breastfed ¹⁶ (%)	41.5
69. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁶ (%)	59.8
70. Breastfeeding children age 6-23 months receiving an adequate diet ^{16, 17} (%)	28.5
71. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{16, 17} (%)	24.9
72. Total children age 6-23 months receiving an adequate diet ^{16, 17} (%)	27.5
73. Children under 5 years who are stunted (height-for-age) ¹⁸ (%)	59.0
74. Children under 5 years who are wasted (weight-for-height) ¹⁸ (%)	8.0
75. Children under 5 years who are severely wasted (weight-for-height) ¹⁹ (%)	2.7
76. Children under 5 years who are underweight (weight-for-age) ¹⁸ (%)	31.1
77. Children under 5 years who are overweight (weight-for-height) ²⁰ (%)	3.1
Nutritional Status of Women (age 15-49 years)	
78. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ²¹ (%)	14.6
79. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ²¹ (%)	8.7
80. Women who have high risk waist-to-hip ratio (≥0.85) (%)	57.8
Anaemia among Children and Women	
81. Children age 6-59 months who are anaemic (<11.0 g/dl) ²² (%)	56.2
82. Non-pregnant women age 15-49 years who are anaemic (<12.0 g/dl) ²² (%)	52.1
83. Pregnant women age 15-49 years who are anaemic (<11.0 g/dl) ²² (%)	49.6
84. All women age 15-49 years who are anaemic ²² (%)	51.8
85. All women age 15-19 years who are anaemic ²² (%)	40.4
Blood Sugar Level among Adults (age 15 years and above)	
Women	
86. Blood sugar level - high (141-160 mg/dl) ²³ (%)	2.3
87. Blood sugar level - very high (>160 mg/dl) ²³ (%)	2.6
88. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ²³ (%)	5.6
Men	
89. Blood sugar level - high (141-160 mg/dl) ²³ (%)	4.3
90. Blood sugar level - very high (>160 mg/dl) ²³ (%)	2.7
91. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ²³ (%)	7.6
Hypertension among Adults (age 15 years and above)	
Women	
92. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	9.1
93. Moderately or severely elevated blood pressure (Systolic ≥160mm of Hg and/or Diastolic ≥100mm of Hg) (%)	4.8
94. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	16.0
Men	
95. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	11.8
96. Moderately or severely elevated blood pressure (Systolic ≥160mm of Hg and/or Diastolic ≥100mm of Hg) (%)	3.8
97. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	16.4
Screening for Cancer among Women (age 30-49 years)	
98. Ever undergone a screening test for cervical cancer (%)	0.8
99. Ever undergone a breast examination for breast cancer (%)	0.0
100. Ever undergone an oral cavity examination for oral cancer (%)	0.2
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)	
101. Women age 15 years and above who use any kind of tobacco (%)	50.2
102. Men age 15 years and above who use any kind of tobacco (%)	73.3
103. Women age 15 years and above who consume alcohol (%)	1.6
104. Men age 15 years and above who consume alcohol (%)	34.4

¹⁵Based on the last child born in the 3 years before the survey.

¹⁶Based on the youngest child living with the mother.

¹⁷Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹⁸Below -2 standard deviations, based on the WHO standard.

¹⁹Below -3 standard deviations, based on the WHO standard.

²⁰Above +2 standard deviations, based on the WHO standard.

²¹Excludes pregnant women and women with a birth in the preceding 2 months.

²²Haemoglobin in grams per decilitre (g/dl). Among children, prevalence is adjusted for altitude. Among women, prevalence is adjusted for altitude and for smoking status, if known. As NFHS uses the capillary blood for estimation of anaemia, the results of NFHS-5 need not be compared with other surveys using venous blood.

²³Random blood sugar measurement.

NOTES

NOTES

INTERNATIONAL INSTITUTE FOR POPULATION SCIENCES

Vision: “To position IIPS as a premier teaching and research institution in population sciences responsive to emerging national and global needs based on values of inclusion, sensitivity and rights protection.”

Mission: “The Institute will strive to be a centre of excellence on population, health and development issues through high quality education, teaching and research. This will be achieved by (a) creating competent professionals, (b) generating and disseminating scientific knowledge and evidence, (c) collaboration and exchange of knowledge, and (d) advocacy and awareness.”

For additional information, please contact:

Director/Principal Investigator (NFHS-5)
International Institute for Population Sciences
Govandi Station Road, Deonar
Mumbai - 400 088 (India)
Telephone: 022 - 42372467
Email: nfhs52017@gmail.com, director@iips.net
Website: <http://www.iipsindia.ac.in>
<http://www.rchiips.org/nfhs/index.shtml>

Director General (Stat.)
Ministry of Health and Family Welfare
Government of India
Indian Red Cross Society Building
Statistics Division
New Delhi 110 001 (India)
Telephone: 011 - 23736979 or 23350003
Email: rajena@nic.in

Chief Director (Stat.)
Ministry of Health and Family Welfare
Government of India
Indian Red Cross Society Building
Statistics Division
New Delhi 110 001 (India)
Telephone: 011 - 23736983
Email: nivedita.g@gov.in
Website: <http://www.mohfw.gov.in>

Technical assistance and additional funding for NFHS-5 was provided by the USAID-supported Demographic and Health Surveys (DHS) Program, ICF, USA. The contents of this publication do not necessarily reflect the views of USAID or the United States Government.



USAID
FROM THE AMERICAN PEOPLE

The opinions in this publication do not necessarily reflect the views of the funding agencies.
For additional information on NFHS-5, visit <http://www.iipsindia.ac.in> or <http://www.mohfw.gov.in>