

Key Indicators for Tripura from NFHS-3	NFHS-3 (2005-06)	Residence		Education				NFHS-2 (1998-99)	NFHS-1 (1992-93)
		Urban	Rural	No education ⁴	<5 years complete ⁴	5-9 years complete ⁴	10 years complete and above ⁴		
Marriage and Fertility									
1. Women age 20-24 married by age 18 (%)	41.6	37.8	42.5	(69.0)	(66.0)	42.2	8.6	37.7	41.1
2. Men age 25-29 married by age 21 (%)	11.4	*	13.0	*	*	(12.3)	(0.0)	na	na
3. Total fertility rate (children per woman)	2.2	1.7	2.3	3.4	2.1	2.0	1.9	1.9	2.7
4. Women age 15-19 who were already mothers or pregnant at the time of the survey (%)	18.5	15.1	19.1	(43.9)	21.8	16.1	6.4	na	na
5. Median age at first birth for women age 25-49	20.3	22.0	20.1	18.8	19.3	20.7	≥25	20.1	19.4
6. Married women with 2 living children wanting no more children ¹ (%)	92.4	91.5	92.6	86.7	92.1	94.3	92.2	87.8	76.3
6a. Two sons	93.8	(89.3)	94.9	(97.2)	*	(92.0)	*	na	na
6b. One son, one daughter	94.4	93.8	94.6	(80.2)	(96.9)	98.4	(90.4)	na	na
6c. Two daughters	84.4	(88.5)	(83.0)	*	*	(86.6)	*	na	na
Family Planning (currently married women, age 15-49)									
Current use									
7. Any method (%)	65.7	66.8	65.5	58.1	72.0	66.0	70.8	55.5	56.4
8. Any modern method (%)	44.9	44.4	45.0	42.5	51.4	44.9	41.0	43.5	28.8
8a. Female sterilization (%)	17.6	16.8	17.8	21.7	21.6	15.6	11.2	26.1	16.9
8b. Male sterilization (%)	0.5	0.3	0.5	1.0	0.9	0.1	0.0	0.6	2.4
8c. IUD (%)	0.9	0.0	1.1	1.0	0.9	0.9	0.6	1.9	1.5
8d. Pill (%)	21.8	23.0	21.6	15.5	25.8	25.2	18.1	13.5	6.4
8e. Condom (%)	3.2	4.3	3.0	0.8	1.7	2.8	10.6	1.4	1.6
Unmet need for family planning									
9. Total unmet need (%)	10.3	8.1	10.7	12.3	8.2	11.0	6.9	17.7	13.5
9a. For spacing (%)	3.8	3.4	3.8	2.5	4.3	4.9	1.8	8.1	5.3
9b. For limiting (%)	6.5	4.7	6.9	9.8	3.9	6.1	5.1	9.6	8.2
Maternal and Child Health									
Maternity care (for births in the last 3 years)									
10. Mothers who had at least 3 antenatal care visits for their last birth (%)	58.7	72.5	56.5	26.6	44.0	69.0	92.9	47.2	38.4
11. Mothers who consumed IFA for 90 days or more when they were pregnant with their last child (%)	18.6	33.3	16.3	9.4	17.6	15.2	42.9	na	na
12. Births assisted by a doctor/nurse/LHV/ANM/other health personnel (%) ²	50.0	79.7	45.4	17.1	27.1	62.2	94.2	47.5	32.2
13. Institutional births (%) ²	48.9	75.7	44.8	16.4	25.9	60.0	96.1	45.2	29.6
14. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/other health personnel within 2 days of delivery for their last birth (%) ²	26.7	30.4	26.1	10.3	11.3	31.9	52.5	na	na
Child immunization and vitamin A supplementation²									
15a. Children 12-23 months fully immunized (BCG, measles, and 3 doses each of polio/DPT) (%)	49.7	*	47.9	(18.7)	*	55.1	*	40.7	19.0
15b. Children 12-23 months who have received BCG (%)	81.1	*	80.2	(61.3)	*	84.0	*	73.5	39.7
15c. Children 12-23 months who have received 3 doses of polio vaccine (%)	65.3	*	64.6	(33.3)	*	70.5	*	57.9	32.2
15d. Children 12-23 months who have received 3 doses of DPT vaccine (%)	60.2	*	59.4	(25.3)	*	66.7	*	51.7	32.2
15e. Children 12-23 months who have received measles vaccine (%)	59.9	*	58.3	(37.3)	*	62.8	*	44.6	28.9
16. Children age 12-35 months who received a vitamin A dose in last 6 months (%)	41.2	(39.6)	41.4	28.0	(36.4)	42.5	(62.4)	na	na
Treatment of childhood diseases (children under 3 years)²									
17. Children with diarrhoea in the last 2 weeks who received ORS (%)	(58.3)	*	(53.1)	*	*	*	*	59.5	41.7
18. Children with diarrhoea in the last 2 weeks taken to a health facility (%)	(69.4)	*	(65.6)	*	*	*	*	62.7	66.7
19. Children with acute respiratory infection or fever in the last 2 weeks taken to a health facility (%)	68.6	*	66.4	(54.4)	*	76.5	*	na	na
Child Feeding Practices and Nutritional Status of Children^{2,3}									
20. Children under 3 years breastfed within one hour of birth (%)	33.1	21.7	35.0	41.2	27.0	34.2	24.4	31.8	6.8
21. Children age 0-5 months exclusively breastfed (%)	36.1	*	38.2	*	*	(30.1)	*	na	na
22. Children age 6-9 months receiving solid or semi-solid food and breastmilk (%)	(59.8)	*	(62.1)	*	*	*	*	na	na
23. Children under 3 years who are stunted (%)	34.1	31.7	34.5	43.6	(36.2)	30.1	30.0	44.6	52.3
24. Children under 3 years who are wasted (%)	24.0	20.0	24.6	34.1	(27.5)	19.0	20.4	18.0	20.6
25. Children under 3 years who are underweight (%)	35.2	25.0	36.7	43.1	(42.0)	30.5	30.7	37.3	42.1
Nutritional Status of Ever-Married Adults (age 15-49)									
26. Women whose Body Mass Index is below normal (%)	35.1	23.8	37.5	42.5	39.9	33.8	16.9	35.2	na
27. Men whose Body Mass Index is below normal (%)	38.3	32.6	39.5	52.0	43.5	38.2	17.5	na	na
28. Women who are overweight or obese (%)	7.8	17.3	5.8	2.6	5.9	7.9	21.2	8.4	na
29. Men who are overweight or obese (%)	5.2	13.5	3.6	0.0	2.0	4.6	16.7	na	na
Anaemia among Children and Adults									
30. Children age 6-35 months who are anaemic (%)	67.9	68.5	67.8	63.3	(65.6)	73.9	(60.3)	61.8	na
31. Ever-married women age 15-49 who are anaemic (%)	67.4	68.9	67.1	70.0	70.2	64.6	67.1	59.0	na
32. Pregnant women age 15-49 who are anaemic (%)	57.6	*	55.6	*	*	(49.5)	*	53.6	na
33. Ever-married men age 15-49 who are anaemic (%)	39.8	41.4	39.5	53.8	37.1	41.8	24.9	na	na
Knowledge of HIV/AIDS among Ever-Married Adults (age 15-49)									
34. Women who have heard of AIDS (%)	72.8	88.2	69.6	42.7	65.9	87.1	99.4	49.0	13.2
35. Men who have heard of AIDS (%)	89.0	98.9	87.0	75.1	84.1	93.7	98.3	na	na
36. Women who know that consistent condom use can reduce the chances of getting HIV/AIDS (%)	42.6	62.4	38.5	19.0	30.1	51.1	82.3	na	na
37. Men who know that consistent condom use can reduce the chances of getting HIV/AIDS (%)	64.7	79.1	61.8	37.9	47.8	74.5	91.4	na	na
Women's Empowerment									
38. Currently married women who usually participate in household decisions (%)	30.2	37.6	28.7	27.4	30.4	30.3	35.1	na	na
39. Ever-married women who have ever experienced spousal violence (%)	44.1	32.2	46.5	56.3	50.2	40.9	20.1	na	na

na : Not available

nc : Not calculated because there are no cases

() Based on 25-49 unweighted cases

* Based on fewer than 25 unweighted cases

¹ Excludes pregnant women.

² Based on the last 2 births in the 3 years before the survey to ever-married women.

³ Based on WHO standard.

⁴ For children education refers to mother's education. Children with missing information on the mother's education are not included in the education columns.