



सत्यमेव जयते  
Government of India

Ministry of Health and Family Welfare

# DISTRICT LEVEL HOUSEHOLD AND FACILITY SURVEY -4

---

## STATE FACT SHEET **MIZORAM**

**(2012-13)**



**IIPS**  
(स्थापना / Established in 1956)  
बेहतर भविष्य के लिए क्षमता निर्माण  
Capacity Building for a Better Future

**International institute for population sciences**  
*(Deemed University)*  
**Mumbai**

## **Introduction**

Three rounds of District Level Household and Facility Survey (DLHS) have been undertaken by the Ministry of Health and Family Welfare, Government of India in the past (Round- I in 1998-99, Round-II in 2002-04, and Round-III in 2007-08) with the main objective to provide reproductive and child health related database at district level in India. The data from these surveys have been useful in setting the benchmarks and examining the progress the country has made after the implementation of RCH programme. In addition, the evidence generated by these surveys have also been useful for the monitoring and evaluation of ongoing programmes and planning of suitable strategies by the central and state governments. In view of the completion of six years of National Rural Health Mission (2005-12), there was a felt need to focus on the achievements and improvements so far. The Ministry of Health and Family Welfare, Government of India, therefore initiated the process of conducting DLHS-4 during 2012-2013 and has designated the International Institute for Population Sciences (IIPS) as the nodal agency to carry out the survey.

Bilingual questionnaires in the local language and English pertaining to Households and Ever Married Women (age 15-49 years) were canvassed by using Computer Assisted Personal Interviewing (CAPI). In DLHS-4 Clinical, Anthropometric and Bio-Chemical (CAB) tests were also carried out in all selected households. The village and health facilities were canvassed using printed questionnaires. In the household questionnaire, information on all members of the household and socio-economic characteristics of the household, assets possessed, number of marriages and deaths in the household since January 2008, etc. were collected. The ever-married women's questionnaire contained information on women's characteristics, maternal care, immunization and childcare, contraception and fertility preferences, reproductive health including knowledge about HIV/AIDS. The village questionnaire contained information on availability of health, education and other facilities in the village, and whether the facilities are accessible throughout the year. The health facility questionnaire contained information on human resources, infrastructure, and services. For the first time, a population-linked facility survey has been conducted in DLHS-4. All Community Health Centres (CHCs), District Hospitals and Sub Divisional Hospitals were covered. Further, all Sub-Health Centres and Primary Health Centres (PHCs) which serve the population of the selected PSUs were also covered. Fieldwork in Mizoram was conducted during June to November 2013, gathering information from 8,583 households, 6,506 ever married women and also from 203 health facilities.

This fact sheet gives information on the key indicators and trends for the state of Mizoram. A separate fact sheet is available for each of the 8 districts.

## Mizoram- Key Indicators

Indicators	DLHS-4 (2012-13)			DLHS-3 (2007-08)		
	TOTAL	RURAL	URBAN	TOTAL	RURAL	URBAN
<b>Sample Size</b>						
Households surveyed.....	8,583	4,986	3,597	10,062	6,456	3,606
Ever married women age 15-49 years.....	6,506	3,820	2,686	7,874	4,991	2,856
Currently married women age 15-49 years interviewed.....	5,676	3,409	2,267	6,845	4,456	2,389
<b>Population and household profile</b>						
Percentage of population literate age 7+ years.....	94.3	90.0	97.9	92.4	89.6	97.4
Percentage of population below age 15 years.....	32.7	36.3	29.5	31.4	32.6	29.2
Mean household size.....	4.5	4.4	4.5	5.1	5.0	5.1
Sex ratio at birth (Males per 100 Females).....	99	102	95	108	113	98
<b>Percentage of households</b>						
Having electricity.....	89.7	78.8	99.5	92.4	88.9	98.7
Improved source of drinking water <sup>1</sup> .....	83.7	71.8	94.4	76.2	68.2	90.5
Having access to improved toilet facility.....	86.4	76.6	95.2	61.3	49.3	83.0
Use clean fuel for cooking <sup>2</sup> .....	52.8	21.4	81.2	36.9	14.8	76.4
<b>Marriage</b>						
Mean age at marriage for girls (marriages that occurred during the reference period).....	23.1	22.2	23.8	21.7	21.2	22.7
Mean age at marriage for boys (marriages that occurred during the reference period).....	26.1	24.6	27.2	24.6	24.3	25.1
Percentage of currently married women married below age 18 years (marriages that occurred during the reference period).....	6.0	7.8	4.7	9.9	11.9	6.3
Percentage of currently married men married below age 21 years (marriages that occurred during the reference period).....	13.5	18.5	9.9	20.9	23.7	16.4
<b>Characteristics of women (%)</b>						
Currently married women who are illiterate.....	10.2	19.7	2.3	9.2	12.8	2.4
Currently married women with 10 or more years of schooling.....	31.5	16.4	44.0	17.9	10.0	32.8
<b>Fertility (%)</b>						
Births to women aged 15-19 years out of total births <sup>3</sup> .....	2.9	3.8	2.2	2.8	3.0	2.3
Women aged 20-24 years reporting birth order of 2 & above.....	46.3	50.2	41.5	39.7	43.6	30.1
Women aged 15-49 years who reported birth order of 3 & above.....	42.1	47.9	36.4	35.0	37.7	29.5
Women with two children wanting no more children.....	10.1	12.3	8.4	13.3	10.4	18.0
Mean no. of children ever born to women age 40-49 years.....	3.4	3.8	3.2	4.0	4.3	3.5
<b>Current use of Family Planning Methods (%)</b>						
Any method.....	59.8	56.9	62.2	53.9	50.6	60.1
Any modern method.....	59.0	56.0	61.5	53.5	50.0	59.6
Female sterilization.....	36.7	33.7	39.1	35.9	33.7	40.2
Male sterilization.....	0.0	0.0	0.0	0.1	0.1	0.2
Pill.....	15.8	17.3	14.5	11.6	11.0	12.8
IUD.....	4.8	3.9	5.5	4.3	4.0	4.9
Condom.....	1.6	0.9	2.1	1.2	1.0	1.5
Any traditional method.....	0.7	0.9	0.7	0.4	0.5	0.3
<b>Unmet Need for Family Planning (%)</b>						
Total unmet need <sup>4</sup> .....	21.4	24.5	18.8	16.7	17.9	14.8
Unmet Need for spacing.....	18.5	20.7	16.6	8.7	8.7	8.8
Unmet Need for limiting.....	2.9	3.8	2.2	6.4	7.1	5.0
Ever use of Emergency Contraceptive Pills (ECP).....	0.1	0.1	0.2	0.8	0.4	1.5

<sup>1</sup> Includes pipe water into dwelling, piped to yard/plot, public tap/standpipe/hand pump/tube well/bore well/well covered/protected spring, tanker/truck, cart with small tank/drum and packaged/bottled water. <sup>2</sup> LPG/PNG/Electricity/Biogas. <sup>3</sup> DLHS-4 reference period is from 1-1-2008 to survey date; DLHS-3 reference period is from 1-1-2004 to survey date.

**Unmet need for spacing** includes fecund women who are neither pregnant nor amenorrhea, who are not using any method of family planning, and say they want to wait two or more years for their next birth. It also includes fecund women who are not using any method of family planning, and say they are unsure whether they want another child or who want another child but are unsure when to have the birth.

**Unmet need for limiting** includes fecund women who are neither pregnant nor amenorrhea, who are not using any method of family planning, and who want no more children. <sup>4</sup>Total unmet need refers to unmet need for limiting and spacing (These definitions are similar to NFHS-3)

## Mizoram - Key Indicators

Indicators	DLHS-4 (2012-13)			DLHS-3 (2007-08)		
	TOTAL	RURAL	URBAN	TOTAL	RURAL	URBAN
<b>Quality of Family Planning Services (%)</b>						
Non-users ever advised by health personnel to adopt any family planning method .....	12.9	12.1	13.6	19.1	19.3	18.7
Current users told about side effects of method .....	13.6	11.3	15.3	27.2	25.7	29.6
Users who received follow-up services for sterilization and IUD within 48 hours .....	27.7	27.0	28.1	50.1	44.4	59.1
Post-partum adoption of family planning for sterilization .....	86.6	84.8	87.9	74.7	73.6	76.3
<b>Antenatal Care (women who had last live/still birth during reference period) (%)</b>						
Pregnant women who received any antenatal check-up.....	83.3	70.0	95.4	89.9	85.6	97.1
Pregnant women who had antenatal check-up in first trimester .....	54.3	40.2	67.3	44.0	37.4	56.6
Pregnant women who had three or more ANC visits .....	61.6	44.6	77.1	62.6	55.8	75.7
Pregnant women who had at least one tetanus toxoid injection .....	80.0	65.6	93.2	86.3	82.2	94.2
Pregnant women whose Blood Pressure (BP) taken .....	73.3	58.2	87.1	68.9	60.9	84.3
Pregnant women who had blood tested (Hb).....	34.7	18.4	49.6	29.0*	18.4*	49.5*
Pregnant women whose abdomen examined .....	50.6	35.5	64.5	66.5	60.3	78.5
Pregnant women who consumed 100 or more IFA Tablets/Syrup equivalent .....	46.5	29.6	62.0	48.4	46.9	51.1
Pregnant women who had full antenatal care <sup>5</sup> .....	36.7	21.0	51.0	33.0	27.4	43.6
<b>Delivery Care (women who had live/still birth during reference period) (%)</b>						
Institutional delivery.....	72.4	49.8	93.0	55.7	40.3	85.6
Delivery at government health institutions.....	62.7	47.1	77.0	50.7	38.8	73.9
Delivery at private health institutions.....	9.7	2.7	16.0	5.0	1.5	11.7
Delivery by Caesarean section at government health institutions.....	3.8	2.4	5.1	2.9	1.8	5.1
Delivery by Caesarean section at private health institutions.....	2.6	0.5	4.6	0.9	0.3	2.1
Delivery at home .....	27.1	49.4	6.7	43.7	59.2	14.0
Delivery at home conducted by skilled health personnel <sup>6</sup> (out of total deliveries).....	7.2	12.4	2.4	7.6	10.3	2.5
Mothers who received post-natal care within 48 hours of Institutional delivery .....	47.1	41.1	50.0	46.9	37.3	64.3
Mothers who received post-natal care within two weeks of Institutional delivery.....	49.7	43.1	53.0	50.5	40.3	68.8
Delivery attended by skilled health personnel.....	79.6	62.2	95.4	63.5	50.7	88.2
Discharge of mothers from institution after minimum stay of 48 hours .....	24.7	19.2	27.5	NA	NA	NA
Out of pocket expenditure per delivery in public health facility (Rs. in 000+).....	2.9	2.9	2.9	NA	NA	NA
<b>Percentage of women who received JSY benefits</b>						
Home delivery .....	3.6	3.3	5.3	4.6	4.1	8.5
Institutional delivery.....	60.9	60.1	61.3	36.8	42.4	31.8
<b>Percentage of Women who had</b>						
Any Pregnancy complication <sup>7</sup> .....	35.8	30.7	40.5	43.9	43.2	45.3
Any Delivery complication <sup>7</sup> .....	10.6	8.0	12.9	19.6	16.8	25.0
Any Post-delivery complication <sup>7</sup> .....	11.0	9.6	12.3	23.9	23.0	25.7
Problem of vaginal discharge during last three months .....	11.9	11.0	12.7	9.9	9.4	10.8
Menstrual related problems during last three months** .....	11.4	10.9	11.8	16.2	16.0	16.5
<b>Percentage of pregnancy resulted in</b>						
Live Birth .....	97.8	97.8	97.9	96.8	97.0	96.4
Still Birth .....	1.1	1.3	0.9	0.5	0.5	0.3
Induced abortion.....	0.1	0.0	0.1	0.1	0.0	0.2
Spontaneous abortion .....	1.0	0.8	1.2	2.7	2.4	3.2
<b>Child Immunization (%) (Children age 12-23 months)</b>						
Number of children.....	705	445	260	697	449	248
Received full vaccination <sup>8</sup> .....	71.2	59.9	82.0	54.5	46.8	68.4
Not received any vaccination .....	6.7	11.8	1.8	4.3	6.0	1.2
Received BCG vaccine .....	88.4	81.3	95.1	92.3	89.2	98.0
Received 3 doses of DPT vaccine .....	82.5	74.3	90.2	66.8	60.9	77.2
Received 3 doses of polio vaccine .....	82.1	72.2	90.9	67.0	59.9	79.9
Received measles vaccine .....	79.7	71.4	87.6	80.4	75.4	89.3
Children (age 9-35 months) received at least one dose of vitamin A supplement in last 6 months .....	57.5	44.2	69.3	72.1	68.0	79.8
<sup>5</sup> Full ANC: At least three visits for antenatal check-up, one TT injection received and 100 IFA tablets or adequate amount of syrup consumed.						
<sup>6</sup> Doctor/ANM/Nurse/midwife/LHV/Other health personnel, <sup>7</sup> Women who had their last live/still birth since 1-1-2008. <sup>8</sup> BCG, 3-injection of DPT, 3 doses of Polio (excluding polio zero) and Measles. Any type of blood tested** Excludes pregnant, in amenorrhea, in menopause, had hysterectomy and ever menstruated women						

## Mizoram - Key Indicators

Indicators	DLHS-4 (2012-13)			DLHS-3 (2007-08)		
	TOTAL	RURAL	URBAN	TOTAL	RURAL	URBAN
<b>Child feeding practices (based on last-born child in the reference period) (%)</b>						
Children age 0-5 months exclusively breastfed <sup>9</sup> .....	69.3	76.4	63.5	59.0	60.6	55.8
Children age 6-9 months receiving solid/semi-solid food and breast milk.....	60.0	55.0	64.5	74.7	70.3	81.0
Children age 12-23 months receiving breast feeding along with complementary feeding**	75.4	74.8	75.9	NA	NA	NA
Children age 6-35 months exclusively breastfed for at least 6 months .....	26.4	25.6	27.2	29.1	28.1	31.0
Children under 3 years breastfed within one hour of birth.....	84.5	81.6	87.0	78.4	77.4	80.1
<b>Birth Weight (%) (age below 36 months)</b>						
Percentage of Children weighed at birth .....	82.9	66.0	98.0	NA	NA	NA
Percentage of Children with low birth weight (out of those who weighed) ( below 2.5 kg) .....	2.9	3.6	2.6	NA	NA	NA
<b>Awareness about Diarrhoea (%)</b>						
Women know about what to do when a child gets diarrhoea.....	71.5	60.1	80.9	85.6	81.5	92.9
<b>Awareness about ARI (%)</b>						
Women aware about danger signs of ARI <sup>10</sup> .....	50.7	37.4	61.7	54.4	46.3	69.7
<b>Treatment of childhood diseases (based on last two surviving children born during the reference period) (%)</b>						
Prevalence of diarrhoea in last 2 weeks for children under 5 year .....	5.5	5.9	5.1	12.2	11.0	14.4
Children with diarrhoea in the last 2 weeks and received ORS <sup>11</sup> .....	60.9	54.5	67.7	55.5	54.7	56.8
Children with diarrhoea in the last 2 weeks given Zinc along with ORS .....	60.1	53.0	67.7	NA	NA	NA
Children with diarrhoea in the last 2 weeks and sought advice/treatment .....	48.4	42.0	55.4	45.0	40.7	51.5
Prevalence of ARI in last 2 weeks for children under 5 years .....	6.6	6.0	7.2	2.1	1.5	3.0
Children with acute respiratory infection or fever in last 2 weeks and sought advice/treatment...	66.7	59.8	72.0	57.5	50.0	67.4
<b>Awareness of RTI/STI and HIV/AIDS (%)</b>						
Women who have heard of RTI/STI.....	31.3	19.9	40.8	45.1	37.3	59.9
Women who have heard of HIV/AIDS.....	84.7	70.0	96.9	92.7	89.3	99.2
Women who have any symptoms of RTI/STI .....	22.6	20.4	24.5	32.6	33.5	30.9
Women who know the place to go for testing of HIV/AIDS <sup>12</sup> .....	74.1	59.2	82.7	59.0	50.5	73.4
Women underwent test for detecting HIV/AIDS <sup>12</sup> .....	51.1	39.1	58.0	9.8	5.9	16.3
<b>Utilization of Government Health Services (%)</b>						
Antenatal care .....	95.4	97.6	94.0	93.2	94.3	91.2
Treatment for pregnancy complications .....	72.7	79.0	70.0	80.1	91.7	65.4
Treatment for post-delivery complications .....	71.1	75.0	68.8	84.4	92.3	74.8
Treatment for vaginal discharge.....	75.2	81.0	72.8	83.4	90.1	73.2
Treatment for children with diarrhoea <sup>13</sup> .....	73.4	66.7	78.8	72.3	78.1	61.2
Treatment for children with ARI <sup>13</sup> .....	58.0	58.1	57.9	58.7	51.9	63.6
<b>Birth Registration (%)</b>						
Children below age 5 years having birth registration done .....	91.2	86.8	95.8	NA	NA	NA
Children below age 5 years who received birth certificate (out of those registered).....	96.9	96.4	97.4	NA	NA	NA
<b>Personal Habits (age 15 years and above) (%)</b>						
Men who use any kind of smokeless tobacco .....	79.3	82.2	76.9	NA	NA	NA
Women who use any kind of smokeless tobacco .....	77.4	79.1	76.0	NA	NA	NA
Men who smoke .....	63.4	68.5	59.0	NA	NA	NA
Women who smoke.....	21.5	26.0	18.0	NA	NA	NA
Men who consume alcohol.....	32.6	33.4	31.9	NA	NA	NA
Women who consume alcohol .....	1.9	2.2	1.8	NA	NA	NA

<sup>9</sup> Children Who were given nothing but breast milk till the survey date <sup>10</sup> Acute Respiratory Infections <sup>11</sup> Oral Rehydration Solutions/Salts. <sup>12</sup> Based on the women who have heard of HIV/AIDS. <sup>13</sup> Last two weeks \*\*includes infant food, tinned powdered or fresh animal milk, fruit juice, tea/coffee or other liquid. na: percentage not shown because of less number of cases.

## Mizoram - Key Indicators

Indicators	DLHS-4 (2012-13)			DLHS-3 (2007-08)		
	TOTAL	RURAL	URBAN	TOTAL	RURAL	URBAN
<b>Reported Prevalence of Morbidity</b>						
Any Injury.....	2.1	2.3	2.0	NA	NA	NA
Acute Illness .....	10.5	10.5	10.5	NA	NA	NA
Chronic Illness .....	3.4	2.8	4.0	NA	NA	NA
<b>Reported Prevalence of Chronic Illness during last one year (%)</b>						
Disease of respiratory system.....	8.8	8.4	9.1	NA	NA	NA
Disease of cardiovascular system .....	4.0	2.5	5.0	NA	NA	NA
Persons suffering from tuberculosis.....	2.8	3.3	2.4	NA	NA	NA
<b>Nutritional status of children below 5 years</b>						
Children below 5 years wasting (weight for height- below 2 SD).....	18.2	19.6	16.0	NA	NA	NA
Children below 5 years wasting (weight for height- below 3 SD).....	6.9	7.5	6.0	NA	NA	NA
Children below 5 years stunting (height for age- below 2 SD).....	37.8	41.1	31.9	NA	NA	NA
Children below 5 years stunting (height for age- below 3 SD).....	15.1	16.3	12.9	NA	NA	NA
Children below 5 years underweight (weight for age- below 2 SD) .....	27.2	31.3	20.3	NA	NA	NA
Children below 5 years underweight (weight for age- below 3 SD) .....	6.9	8.5	4.3	NA	NA	NA
<b>Anaemia Status by Haemoglobin Level<sup>14</sup> (%)</b>						
Children (6-59 months) having anaemia.....	70.7	72.1	69.4	NA	NA	NA
Children (6-59 months) having severe anaemia.....	13.0	14.6	11.5	NA	NA	NA
Children (6-9 Years) having anaemia - Male .....	69.1	71.6	66.2	NA	NA	NA
Children (6-9 Years) having severe anaemia - Male .....	11.2	13.3	8.7	NA	NA	NA
Children (6-9 Years) having anaemia - Female.....	70.5	73.4	67.2	NA	NA	NA
Children (6-9 Years) having severe anaemia - Female .....	12.6	14.8	10.1	NA	NA	NA
Children (6-14 years) having anaemia - Male.....	67.1	69.6	64.6	NA	NA	NA
Children (6-14 years) having severe anaemia - Male.....	10.5	12.8	7.9	NA	NA	NA
Children (6-14 years) having anaemia - Female .....	67.7	69.1	66.2	NA	NA	NA
Children (6-14 years) having severe anaemia - Female .....	10.6	11.7	9.3	NA	NA	NA
Children (10-19 Years <sup>15</sup> ) having anaemia - Male.....	61.1	65.1	57.1	NA	NA	NA
Children (10-19 Years <sup>15</sup> ) having severe anaemia - Male.....	9.1	11.3	7.0	NA	NA	NA
Children (10-19 Years <sup>15</sup> ) having anaemia - Female .....	64.4	66.0	62.9	NA	NA	NA
Children (10-19 Years <sup>15</sup> ) having severe anaemia - Female.....	9.0	9.9	8.0	NA	NA	NA
Adolescents (15-19 years) having anaemia .....	58.8	63.7	55.0	NA	NA	NA
Adolescents (15-19 years) having severe anaemia .....	8.2	10.4	6.5	NA	NA	NA
Pregnant women (15-49 aged) having anaemia .....	63.2	64.4	62.1	NA	NA	NA
Pregnant women (15-49 aged) having severe anaemia.....	8.4	9.8	7.1	NA	NA	NA
Women (15-49 aged) having anaemia .....	64.1	66.1	62.4	NA	NA	NA
Women (15-49 aged) having severe anaemia.....	8.4	9.1	7.7	NA	NA	NA
Persons (20 years and above) having anaemia .....	55.7	59.3	52.6	NA	NA	NA
Persons (20 years and above) having Severe anaemia.....	6.6	7.5	5.9	NA	NA	NA
<b>Blood Sugar Level (age 18 years and above) (%)</b>						
Blood Sugar Level >140 mg/dl (high) .....	8.4	7.6	9.0	NA	NA	NA
Blood Sugar Level >160 mg/dl (very high) .....	4.1	3.4	4.6	NA	NA	NA
<b>Hypertension (age 18 years and above) (%)</b>						
Above Normal Range (Systolic >140 mm of Hg & Diastolic >90 mm of Hg ) .....	19.8	18.1	21.1	NA	NA	NA
Moderately High (Systolic >160 mm of Hg & Diastolic >100 mm of Hg ) .....	5.4	4.2	6.4	NA	NA	NA
Very High (Systolic >180 mm of Hg & Diastolic >110 mm of Hg ).....	1.8	1.4	2.1	NA	NA	NA
<sup>14</sup> Any anaemia below 11g/dl, severe anaemia below 7g/dl. <sup>15</sup> Excluding age group 19 years						
<b>Chronic Illness</b> :Any person with symptoms persisting for longer than one month is defined as suffering from chronic illness						

## Mizoram- Key Indicators

Indicators	Number/Percentage	
	DLHS-4	DLHS-3
<b>Villages covered</b>		
Number of villages.....	184	245
<b>Health Facilities covered</b>		
Number of Sub-Health Centres.....	140	172
Number of Primary Health Centres (PHC) .....	42	53
Number of Community Health Centres (CHC) including Block PHC .....	11	10
Number of Sub-Divisional Hospitals (SDH) .....	02	NA
Number of District Hospitals (DH) .....	08	08
<b>Health programmes at village level</b>		
Percentage of villages having ASHA.....	84.8	69.4
Percentage of Villages having Village Health Nutrition and Sanitation Committee (VHNSC).....	69.0	88.2
<b>Accessibility of health facility (%)</b>		
Villages with Sub-Health Centre within 3 km .....	72.8	69.4
Villages with PHC within 10 km .....	39.8	28.6
<b>Availability of Health Infrastructure, Staff and Services (%)</b>		
<b>Sub-Health Centre</b>		
Sub-Health Centre located in government building .....	98.6	97.1
Sub-Health Centre with ANM.....	89.2	95.4
Sub-Health Centre with male health worker .....	64.3	93.0
Sub-Health Centre with ANM residing in Sub-Health Centre quarter where facility is available .....	76.6	79.1
Sub-Health Centre with additional ANM .....	13.8	38.4
<b>Primary Health Centre (PHC)</b>		
PHCs functioning on 24 X 7 hours basis .....	90.2	69.8
PHCs having Lady Medical Officer *.....	12.8	22.6
PHCs with at least 4 beds .....	95.2	92.5
PHCs with AYUSH doctor* .....	0	1.9
PHCs having residential quarter for Medical Officer .....	95.2	88.7
PHCs having new born care services on 24 X 7 hours basis .....	92.3	91.9
PHCs having referral services for pregnancies/delivery on 24 X 7 hours basis .....	16.2	29.7
PHCs conducted at least 10 deliveries during last one month on 24 X 7 hours basis.....	10.3	29.7
<b>Community Health Centre (CHC)</b>		
CHCs having 24 X 7 hours normal delivery services .....	100	90.0
CHCs having Obstetrician/Gynaecologist .....	18.2	0.0
CHCs having Anesthetist.....	27.3	10.0
CHCs having functional Operation Theatre .....	100	80.0
CHCs designated as FRUs.....	54.6	70.0
CHCs designated as FRUs offering caesarean section .....	100.0	0.0
CHCs having new born care services on 24 X 7 hours basis .....	100.0	100.0
<b>Sub Divisional Hospital (SDH)</b>		
SDHs having Paediatrician .....	50.0	NA
SDHs having regular radiographer .....	0.0	NA
SDHs having 2D Echo facility .....	0.0	NA
SDHs having Ultrasound facility .....	0.0	NA
SDHs having three phase connection .....	0.0	NA
SDHs having critical care area .....	0.0	NA
SDHs having suggestion and complaint box .....	100.0	NA
<b>District Hospital (DH)</b>		
DHs having Paediatrician .....	50.0	NA
DHs having regular radiographer.....	12.5	NA
DHs having 2D Echo facility .....	25.0	NA
DHs having Ultrasound facility.....	100.0	NA
DHs having three phase connection.....	62.5	NA
DHs having critical care area.....	35.5	NA
DHs having suggestion and complaint box.....	87.5	NA

\* Out of total medical officers available

# INTERNATIONAL INSTITUTE FOR POPULATION SCIENCES

**Vision:** “To position IIPS as a premier teaching and research institution in population sciences responsive to emerging national and global needs based on values of inclusion, sensitivity and rights protection.”

**Mission:** “The Institute will strive to be a centre of excellence on population, health and development issues through high quality education, teaching and research. This will be achieved by (a) creating competent professionals, (b) generating and disseminating scientific knowledge and evidence, (c) collaboration and exchange of knowledge, and (d) advocacy and awareness.”

## **For additional information, please contact:**

### **Director/Project Coordinator (DLHS-4) International Institute for Population Sciences**

Govandi Station Road, Deonar  
Mumbai - 400 088 (India)  
Telephone: 022-4237 2465, 42372411  
Fax: 022-25563257, 25555895  
Email: rchpro@iips.net, director@iips.net  
Website: <http://www.rchiips.org>  
<http://www.iipsindia.org>

### **Additional Director General (Stat.) Ministry of Health and Family Welfare Government of India**

Nirman Bhavan  
New Chandigarh 110 011  
Telephone: 011 - 23061334  
Fax: 011 - 23061334  
Email: adg-mohfw@nic.in

### **Chief Director (Stat.) Ministry of Health and Family Welfare Government of India**

Nirman Bhavan  
New Chandigarh 110 011  
Telephone: 011 - 23062699  
Fax: 011 - 23062699  
Email: cdstat@nic.in  
Website: <http://www.mohfw.nic.in>